

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Roxatidine Acetate (hydrochloride)

Item No. 21248

CAS Registry No.:	93793-83-0				
Formal Name:	2-(acetyloxy)-N-[3-[3-(1-piperidinylmethyl)				
	phenoxy]propyl]-acetamide, monohydrochloride				
Synonyms:	HOE 760, TZU-0460	o H			
MF:	$C_{19}H_{28}N_2O_4 \bullet HCI$	Ŭ ~ N	0.	\sim \sim	\sim
FW:	384.9		\sim \sim \sim	\downarrow	`N \
Purity:	≥98%	Ö			
UV/Vis.:	λ _{max} : 223, 277 nm		• HCI	\checkmark	\searrow
Supplied as:	A crystalline solid				
Storage:	-20°C				
Stability:	≥2 years				
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.					

Laboratory Procedures

Roxatidine acetate (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the roxatidine acetate (hydrochloride) in the solvent of choice. Roxatidine acetate (hydrochloride) is soluble in organic solvents such as ethanol and DMSO. It is also soluble in water. The solubility of roxatidine acetate (hydrochloride) in ethanol, DMSO, and water is approximately 12, 78, and 77 mg/ml, respectively. We do not recommend storing the aqueous solution for more than one day.

Description

Roxatidine acetate is a competitive histamine H₂ receptor antagonist that upon oral absorption is rapidly converted to its active metabolite roxatidine.¹ By inhibiting the binding of histamine to H₂ receptors, roxatidine reduces both intracellular cAMP concentrations and gastric acid secretion by parietal cells.¹

Reference

1. Murdoch, D., and McTavish, D. Roxatidine acetate. A review of its pharmacodynamic and pharmacokinetic properties, and its therapeutic potential in peptic ulcer disease and related disorders. Drugs 42(2), 240-260 (1991).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Super agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/17/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM