

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



DCP-LA

Item No. 21375

CAS Registry No.: 28399-31-7

Formal Name: 2-[(2-pentylcyclopropyl)methyl]-

cyclopropaneoctanoic acid

Synonym: FR 236924 MF: $C_{20}H_{36}O_{2}$ FW: 308.5

Purity: ≥95%

Supplied as: A solution in ethanol

Storage: -20°C Stability: ≥1 year

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

DCP-LA is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol, DMSO, and dimethyl formamide purged with an inert gas can be used. The solubility of DCP-LA in these solvents is approximately 10, 25, and 50 mg/ml, respectively.

Description

DCP-LA is a derivative of linoleic acid (Item No. 90150) and a potent activator of PKC_E.¹ It activates PKC ϵ with a greater than 7-fold stronger potency over PKC α , βI , βII , γ , δ , μ , η , and ζ in a cell-free assay. DCP-LA activates PKC in PC12 cells in a concentration-dependent manner, an effect that is blocked by the PCK inhibitor bisindolylmaleimide I (Item Nos. 13298 | 21180) and a PCKε-selective peptide inhibitor. It decreases intracellular levels of amyloid-β (Aβ) in Neuro2 neuroblastoma cells transfected with human APPSwe/PS $1\Delta9.^2$ DCP-LA also stimulates hippocampal glutamate release, striatal dopamine release, and hypothalamic serotonin release in rat brain slices in a PKC- and α7-containing nicotinic acetylcholine receptor-dependent manner.³ In vivo, DCP-LA (3 mg/kg, i.p.) prevents synaptic loss and amyloid plaque formation and decreases escape latency in the Morris water maze in the 5XFAD transgenic mouse model of Alzheimer's disease.4

References

- 1. Kanno, T., Yamamoto, H., Yaguchi, T., et al. The linoleic acid derivative DCP-LA selectively activates PKC-ε, possibly binding to the phosphatidylserine binding site. J. Lipid. Res. 47(6), 1146-1156 (2006).
- 2. Nelson, T.J., Cui, C., Luo, Y., et al. Reduction of β-amyloid levels by novel protein kinase Cε activators. J. Biol. Chem. 284(50), 34514-34521 (2009).
- 3. Shimizu, T., Kanno, T., Tanaka, A., et al. α, β-DCP-LA selectively activates PKC-ε and stimulates neurotransmitter release with the highest potency among 4 diastereomers. Cell Physiol. Biochem. 27(2), 149-158 (2011).
- 4. Hongpaisan, J., Sun, M.-K., and Alkon, D.L. PKC ε activation prevents synaptic loss, Aβ elevation, and cognitive deficits in Alzheimer's disease transgenic mice. J. Neurosci. 31(2), 630-643 (2011).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/11/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM