

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



(R,S)-CHPG

Item No. 21408

CAS Registry No.: 170846-74-9

Formal Name: α-amino-2-chloro-5-hydroxy-benzeneacetic acid

Synonyms: Chlorohydroxyphenylglycine, CHPG

MF: C₈H₈CINO₃ FW: 201.6 **Purity:** ≥98% Supplied as: A solid Storage: -20°C Stability: ≥2 years

Special Conditions: Store in desiccating conditions

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

(R,S)-CHPG is supplied as a solid. A stock solution may be made by dissolving the (R,S)-CHPG in the solvent of choice. (R,S)-CHPG is soluble in 1 eq. NaOH to 100 mM.

Description

(R,S)-CHPG is a potent and selective agonist at metabotropic glutamate receptor 5 (mGluR5). In CHO cells, it had an EC₅₀ of 750 μM for calcium mobilization in mGluR5a-expressing cells but was inactive in mGluR1a-expressing cells. In transfected HEK293 cells, (R,S)-CHPG bound mGluR1a and mGluR5a $(K_s = 0.9 \text{ and } 3.9 \mu\text{M}, \text{ respectively})$ but not ionotropic glutamate receptors.² It reduced oxidative stress and inflammatory markers in cultured BV-2 microglial cells.³ In mouse models of traumatic brain injury, it reduced lesion volume, improved sensorimotor deficits in the beam walk test, and improved spatial memory in the Morris water maze.4

References

- 1. Doherty, A.J., Palmer, M.J., Henley, J.M., et al. (RS)-2-chloro-5-hydroxyphenylglycine (CHPG) activates mGlu₅, but no mGlu₁, receptors expressed in CHO cells and potentiates NMDA responses in the hippocampus. Neuropharmacology 36(2), 265-267 (1997).
- 2. Mutel, V., Ellis, G.J., Adam, G., et al. Characterization of [3H]quisqualate binding to recombinant rat metabotropic glutamate 1a and 5a receptors and to rat and human brain sections. J. Neurochem. 75(6), 2590-2601 (2000).
- 3. Qiu, J.-L., Zhu, W.-L., Lu, Y.-J., et al. The selective mGluR5 agonist CHPG attenuates SO₂-induced oxidative stress and inflammation through TSG-6/NF-kB pathway in BV2 microglial cells. Neurochem Int. **85-86**, 46-52 (2015).
- 4. Loane, D.J., Stoica, B.A., Byrnes, K.R., et al. Activation of mGluR5 and inhibition of NADPH oxidase improves functional recovery after traumatic brain injury. J. Neurotrauma 30(5), 403-412 (2013).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 10/24/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM