

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Cisapride

Item No. 21657

CAS Registry No.: 81098-60-4

Formal Name: rel-4-amino-5-chloro-N-[1-

> [(3R,4S)-3-(4-fluorophenoxy) propyl]-3-methoxy-4-piperidinyl]-

2-methoxy-benzamide

C₂₃H₂₉CIFN₃O₄ MF:


FW: 466.0 **Purity:**

UV/Vis.: λ_{max}: 214, 276, 308 nm

Supplied as: A crystalline solid

Storage: Stability:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

Cisapride is supplied as a crystalline solid. A stock solution may be made by dissolving the cisapride in the solvent of choice, which should be purged with an inert gas. Cisapride is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of cisapride in these solvents is approximately 30 mg/ml. Cisapride is also slightly soluble in ethanol.

Cisapride is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, cisapride should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Cisapride has a solubility of approximately 0.2 mg/ml in a 1:4 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Cisapride is an agonist of serotonin (5-HT) receptor subtype 5-HT₄ (IC₅₀ = $0.483 \mu M$ in COS-7 cells expressing the human receptor). It induces relaxation of precontracted isolated rat esophageal thoracic muscularis mucosae preparations (EC₅₀ = 102.33 nM). Cisapride (0.5 mg/kg) increases the rate of gastric emptying in rats. It is also a human ether-a-go-go related gene (hERG) channel blocker that binds to hERG channels with an IC₅₀ value of less than 1 μ M in a fluorescence polarization assay. Formulations containing cisapride have previously been used in the treatment of nocturnal heartburn associated with gastroesophageal reflux disease.

Reference

1. Park, J.S., Im, W., Choi, S., et al. Discovery and SAR of N-(1-((substituted piperidin-4-yl)methyl)-3methoxypiperidin-4-yl)-2-methoxybenzamide derivatives: 5-Hydroxytryptamine receptor 4 agonist as a potent prokinetic agent. Eur. J. Med. Chem. 109, 75-88 (2016).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 12/03/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM