



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



P1075

Item No. 21849

CAS Registry No.: 60559-98-0
Formal Name: N-cyano-N'-(1,1-dimethylpropyl)-N''-3-pyridinyl-guanidine
MF: C₁₂H₁₇N₅
FW: 231.3
Purity: ≥98%
UV/Vis.: λ_{max}: 239 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

P1075 is supplied as a crystalline solid. A stock solution may be made by dissolving the P1075 in the solvent of choice. P1075 is soluble in organic solvents such as ethanol and DMSO, which should be purged with an inert gas. The solubility of P1075 in these solvents is up to approximately 50 and 100 mM, respectively.

Description

P1075 is a potent activator of sulfonylurea receptor 2-associated ATP-sensitive potassium channels (SUR2-K_{ir}6) with an EC₅₀ value of 45 nM for SUR2B-K_{ir}6 channel activation.¹ It is highly selective for SUR2 over SUR1 isoforms (IC₅₀s = 9-46 nM and 1.02 mM, respectively, in a radioligand binding assay). P1075 increases outflow facility, a marker of reduced intraocular pressure, in human eye anterior segments *ex vivo*.² It also reduces infarct size in isolated rabbit hearts *via* activation of mitochondrial ATP-sensitive potassium channels (K_{ATP}; EC₅₀s = 60-90 nM).³

References

- Schwanstecher, M., Sieverding, C., Dörschner, H., *et al.* Potassium channel openers require ATP to bind to and act through sulfonylurea receptors. *EMBO J.* **17(19)**, 5529-5535 (1998).
- Chowdhury, U.R., Bahler, C.K., Hann, C.R., *et al.* ATP-sensitive potassium (K_{ATP}) channel activation decreases intraocular pressure in the anterior chamber of the eye. *Invest. Ophthalmol. Vis. Sci.* **52(9)**, 6435-6442 (2011).
- Oldenburg, O., Yang, X.M., Krieg, T., *et al.* P1075 opens mitochondrial K_{ATP} channels and generates reactive oxygen species resulting in cardioprotection of rabbit hearts. *J. Mol. Cell. Cardiol.* **35(9)**, 1035-1042 (2003).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/15/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM