

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



EGA

Item No. 21882

| CAS Registry No.:  | 415687-81-9  |      |
|--|--|------|
| Formal Name:   | 2-[(4-bromophenyl)methylene]-                      |      |
|  | N-(2,6-dimethylphenyl)-<br>hydrazinecarboxamide    |      |
| MF:  | C <sub>16</sub> H <sub>16</sub> BrN <sub>3</sub> O | H Br |
| FW:  | 346.2  |      |
| Purity:  | ≥98%   |      |
| UV/Vis.:   | λ <sub>max</sub> : 291 nm                          |      |
| Supplied as:   | A crystalline solid                                | H    |
| Storage:   | -20°C  |      |
| Stability:   | ≥2 years   |      |
| Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis. |  |      |

#### Laboratory Procedures

EGA is supplied as a crystalline solid. A stock solution may be made by dissolving the EGA in the solvent of choice. EGA is soluble in the organic solvent DMSO, which should be purged with an inert gas, at a concentration of approximately 5 mg/ml.

#### Description

EGA is an inhibitor of endosomal trafficking.<sup>1</sup> It increases accumulation of fluorescently labeled EGF in early endosomes in HeLa cells when used at a concentration of 20 μM. EGA inhibits cell death induced by anthrax lethal toxin in RAW264.7 cells (IC<sub>50</sub> = 1.7  $\mu$ M) as well as by additional acid-dependent bacterial toxins, including diphtheria toxin, P. aeruginosa ExoA, and H. ducreyi cytolethal distending toxin (Hd-CDT), in various cell types. It also inhibits infection of Vero and HeLa cells by the low pH-dependent lymphocytic choriomeningitis (LCMV; Armstrong strain) and influenza A/WSN/33 viruses, respectively.

#### Reference

1. Gillespie, E.J., Ho, C.L., Balaji, K., et al. Selective inhibitor of endosomal trafficking pathways exploited by multiple toxins and viruses. Proc. Natl. Acad. Sci. U.S.A. 110(50), E4904-E4912 (2013).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Super agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/31/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM