

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Linoleic Acid (sodium salt)

Item No. 21909

CAS Registry No.: 822-17-3

Formal Name: 9Z,12Z-octadecadienoic acid,

monosodium salt

Synonym: Telfairic Acid MF: C₁₈H₃₁O₂ • Na

FW: 302.4 **Purity:** ≥98%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Linoleic acid (sodium salt) is supplied as a crystalline solid. A stock solution may be made by dissolving the linoleic acid (sodium salt) in the solvent of choice. Linoleic acid (sodium salt) is soluble in the organic solvent ethanol, which should be purged with an inert gas, at a concentration of approximately 1.5 mg/ml.

Linoleic acid (sodium salt) is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, linoleic acid (sodium salt) should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Linoleic acid (sodium salt) has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol: PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Linoleic acid is an essential ω -6 polyunsaturated fatty acid (PUFA). It is the most abundant PUFA in a variety of foods, and dietary sources of linoleic acid include vegetable oils, meats, nuts, seeds, and eggs. Linoleic acid (30 μM) increases migration of IEC-6 rat intestinal epithelial cells in a wound healing assay.² Rats fed a linoleate-deficient diet exhibit decreased body weight and an increased ratio of eicosatrienoate to arachidonate in liver and serum phospholipids compared with rats fed a control diet, as well as mild scaling of forepaw skin.3

References

- 1. Whelan, J. and Fritsche, K. Linoleic acid. Adv. Nutr. 4(3), 311-312 (2013).
- 2. Ruthig, D.J., and Meckling-Gill, K.A. Both (n-3) and (n-6) fatty acids stimulate wound healing in the rat intestinal epithelial cell line, IEC-6. J. Nutr. 129(10), 1791-1798 (1999).
- 3. Cunnane, S.C. and Anderson, M.J. Pure linoleate deficiency in the rat: Influence on growth, accumulation of n-6 polyunsaturates, and [1-14C]linoleate oxidation. J. Lipid Res. 38(4), 805-812 (1997).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/08/2020

COO

• Nat

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM