



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



MONNA

Item No. 21917

CAS Registry No.: 1572936-83-4
Formal Name: 2-[[4-methoxy-2-naphthalenyl) amino]-5-nitro-benzoic acid
MF: C₁₈H₁₄N₂O₅
FW: 338.3
Purity: ≥98%
UV/Vis.: λ_{max}: 214, 264, 295, 376 nm
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

MONNA is supplied as a crystalline solid. A stock solution may be made by dissolving the MONNA in the solvent of choice. MONNA is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of MONNA in these solvents is approximately 30 mg/ml. MONNA is also slightly soluble in ethanol.

MONNA is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, MONNA should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. MONNA has a solubility of approximately 0.25 mg/ml in a 1:3 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

MONNA is a selective inhibitor of anoctamin-1 (ANO1; IC₅₀ = 0.08 μM in *Xenopus* oocytes), a calcium-activated chloride channel (CaCC).¹ It is selective for ANO1 over the chloride channels bestrophin-1, chloride channel protein 2, and cystic fibrosis transmembrane conductance regulator up to 10 μM. MONNA (10 μM) induces full vasorelaxation of precontracted mouse isolated mesenteric arteries in the presence or absence of chloride.² It also hyperpolarizes isolated rat mesenteric arteries under resting conditions.

References

1. Oh, S. J., Hwang, S.J., Jung, J.E., *et al.* MONNA, a potent and selective blocker for transmembrane protein with unknown function 16/anoctamin-1. *Mol. Pharmacol.* **84**(5), 726-735 (2013).
2. Boedtjker, D.M.B., Kim, S., Jensen, A.B., *et al.* New selective inhibitors of calcium-activated chloride channels - T16A_{inh}-A01, CaCC_{inh}-A01 and MONNA - what do they inhibit? *Br. J. Pharmacol.* **172**(16), 4158-4172 (2015).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/02/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM