

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



Dexamethasone-d₄

Item No. 22100

CAS Registry No.:	2483831-63-4	
Formal Name:	9-fluoro-11β,17,21-trihydroxy-16α-methyl-	
	pregna-1,4-diene-3,20-dione-4,6,21,21-d ₄	
MF:	$C_{22}H_{25}D_4FO_5$	
FW:	396.5	HO
Chemical Purity:	≥98% (Dexamethasone)	
Deuterium		
Incorporation:	≥99% deuterated forms (d₁-d₄); ≤1% d₀	F F
Supplied as:	A solid	
Storage:	-20°C	
Stability:	≥2 years	
Special Conditions: Heat and light sensitive		

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Dexame thas one- d_4 is intended for use as an internal standard for the quantification of dexame thas one (Item No. 11015) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Dexamethasone- d_4 is supplied as a solid. A stock solution may be made by dissolving the dexamethasone- d_4 in the solvent of choice. Dexamethasone- d_4 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of dexamethasone- d_4 in these solvents is approximately 3, 30, and 25 mg/ml, respectively.

Description

Dexamethasone is a synthetic glucocorticoid that binds the human glucocorticoid receptor with a higher affinity than a natural ligand, cortisol (K_d = 5 nM versus 17 nM, respectively).¹ Through receptor activation, dexamethasone has both transactivating and transrepressing effects on gene expression, producing, in general, anti-inflammatory results.²

References

- 1. Mulatero, P., Panarelli, M., Schiavone, D., et al. Impaired cortisol binding to glucocorticoid receptors in hypertensive patients. Hypertension 30(5), 1274-1278 (1997).
- 2. Beck, I.M., Vanden Berghe, W., Vermeulen, L., et al. Crosstalk in inflammation: The interplay of glucocorticoid receptor-based mechanisms and kinases and phosphatases. Endocr. Rev. 30(7), 830-882 (2009).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 12/21/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM