



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Fedotozine (tartrate)

Item No. 22189

CAS Registry No.: 133267-27-3
Formal Name: (R)- α -ethyl-N,N-dimethyl- α -[[[(3,4,5-trimethoxyphenyl)methoxy]methyl]-benzenemethanamine, [S-(R*,R*)]-2,3-dihydroxybutanedioate
MF: C₂₂H₃₁NO₄ • C₄H₆O₆
FW: 523.6
Purity: ≥95%
Supplied as: A solid
Storage: -20°C
Stability: ≥4 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Fedotozine (tartrate) is supplied as a solid. A stock solution may be made by dissolving the fedotozine (tartrate) in the solvent of choice, which should be purged with an inert gas. Fedotozine (tartrate) is sparingly soluble (1-10 mg/ml) in DMSO.

Fedotozine (tartrate) is slightly soluble (0.1-1 mg/ml) in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

Description

Fedotozine is a κ -opioid receptor agonist.¹ It binds to κ -opioid receptors in mouse brain membranes ($K_i = 0.16$ nM). *In vivo*, fedotozine (10 mg/kg) reduces gastric extension-induced increases in mean arterial pressure (MAP), a marker for hyperalgesia, in a rat model of duodenal acidification-induced gastric hypersensitivity.² It also prevents colonic distension-induced inhibition of gastric motility and emptying in dogs when administered at a dose of 50 μ g/kg.³

References

1. Lai, J., Ma, S.W., Zhu, R.H., *et al.* Pharmacological characterization of the cloned κ opioid receptor as a κ_{1b} subtype. *Neuroreport* **5(16)**, 2161-2164 (1994).
2. Nakata-Fukuda, M., Hirata, T., Keto, Y., *et al.* Inhibitory effect of the selective serotonin 5-HT₃ receptor antagonist ramosetron on duodenal acidification-induced gastric hypersensitivity in rats. *Eur. J. Pharmacol.* **731**, 88-92 (2014).
3. Gué, M., Junien, J.L., and Buéno, L. The κ agonist fedotozine modulates colonic distention-induced inhibition of gastric motility and emptying in dogs. *Gastroenterol.* **107(5)**, 1327-1334 (1994).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/09/2024

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM