

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Melatonin-d₄

Item No. 22291

CAS Registry No.: 66521-38-8

Formal Name: N-[2-(5-methoxy-1H-indol-3-yl)ethyl-

 $1,1,2,2-d_{4}$]-acetamide

MF: $C_{13}H_{12}D_4N_2O_2$

FW: 236.3

Chemical Purity:

≥98% (Melatonin)

Deuterium

Incorporation: ≥99% deuterated forms (d1-d4); ≤1% d0

 λ_{max} : 225, 280 nm A crystalline solid UV/Vis.: Supplied as:

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Melatonin-d₄ is intended for use as an internal standard for the quantification of melatonin (Item No. 14427) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

 $Melatonin-d_4 is supplied as a crystalline solid. A stock solution may be made by dissolving the melatonin-d_4$ in the solvent of choice. Melatonin- d_4 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of melatonin- d_a in ethanol is approximately 20 mg/ml and approximately 30 mg/ml in DMSO and DMF.

Description

Melatonin is an indoleamine neurohormone whose levels vary in a daily cycle, thereby allowing the entrainment of the circadian rhythms of several biological functions in animals, plants, and microbes.¹ Many biological effects of melatonin are transduced through melatonin receptors, including the MT₁, MT₂, and MT₂ subtypes. Melatonin also acts as a powerful antioxidant that protects lipids, proteins, and DNA against oxidative damage.^{2,3} Glutathione peroxidase, superoxide dismutases, and catalase are upregulated by melatonin, and melatonin scavenges free radicals as a terminal antioxidant.^{2,3}

References

- 1. Boutin, J.A., Audinot, V., Ferry, G., et al. Molecular tools to study melatonin pathways and actions. Trends Pharamacol. Sci. 26(8), 412-419 (2005).
- Reiter, R.J., Tan, D.X., Manchester, L.C., et al. Biochemical reactivity of melatonin with reactive oxygen and nitrogen species. Cell Biochem. Biophys. 34(2), 237-256 (2001).
- Reiter, R.J., Tang, L., Garcia, J.J., et al. Pharmacological actions of melatonin in oxygen radical pathophysiology. Life Sci. 60(25), 2255-2271 (1997).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/25/2017

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM