



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Sphinganine (d17:0)

Item No. 22510

CAS Registry No.: 32164-02-6
Formal Name: 2S-amino-1,3R-heptadecanediol
Synonyms: D-erythro-C17-Dihydrosphingosine,
D-erythro-Sphinganine C-17
MF: C₁₇H₃₇NO₂
FW: 287.5
Purity: ≥95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Sphinganine (d17:0) is supplied as a crystalline solid. A stock solution may be made by dissolving the sphinganine (d17:0) in the solvent of choice. Sphinganine (d17:0) is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of sphinganine (d17:0) in these solvents is approximately 2 and 10 mg/ml, respectively. Sphinganine (d17:0) is also miscible in ethanol.

Description

Sphinganine is a synthetic bioactive sphingolipid that inhibits the growth of *C. glabrata* and *C. albicans* with a minimum fungicidal concentration (MFC) value of 0.5 µg/ml for both.^{1,2} More commonly, sphinganine is used as an internal standard in the analysis of sphingoid compounds by chromatographic or spectrometric methods.^{2,3}

References

1. Thevissen, K., Hillaert, U., Meert, E.M., *et al.* Fungicidal activity of truncated analogues of dihydrosphingosine. *Bioorg. Med. Chem. Lett.* **18(13)**, 3728-3730 (2008).
2. Spassieva, S., Bielawski, J., Anelli, V., *et al.* Combination of C17 sphingoid base homologues and mass spectrometry analysis as a new approach to study sphingolipid metabolism. *Methods Enzymol.* **434**, 233-241 (2007).
3. Qu, F., Wu, C.S., Hou, J.F., *et al.* Sphingolipids as new biomarkers for assessment of delayed-type hypersensitivity and response to triptolide. *PLoS One* **7(12)**, e52454 (2012).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/08/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM