



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## C22 Ceramide (d18:1/22:0)

Item No. 22533

**CAS Registry No.:** 27888-44-4  
**Formal Name:** N-[(1S,2R,3E)-2-hydroxy-1-(hydroxymethyl)-3-heptadecen-1-yl]-docosanamide  
**Synonyms:** C22 Ceramide, Cer(d18:1/22:0), Ceramide (d18:1/22:0)  
**MF:** C<sub>40</sub>H<sub>79</sub>NO<sub>3</sub>  
**FW:** 622.1  
**Purity:** ≥95%  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

C22 Ceramide (d18:1/22:0) is supplied as a crystalline solid. A stock solution may be made by dissolving the C22 ceramide (d18:1/22:0) in the solvent of choice, which should be purged with an inert gas. C22 Ceramide (d18:1/22:0) is soluble in the organic solvent dimethyl formamide at a concentration of approximately 0.15 mg/ml.

### Description

C22 Ceramide is an endogenous bioactive sphingolipid. Ceramides are involved in permeabilization of the mitochondrial outer membrane during apoptosis via the intrinsic pathway.<sup>1</sup> C22 Ceramide forms small channels in liposomes, whereas C16 ceramide (Item No. 10681) forms channels that can grow in size, suggesting that acyl chain length of ceramides is important in mitochondrial-mediated apoptosis. In addition, the mitochondrial-to-cytosolic stress response (MCSR) in *C. elegans* fed C22 ceramide decreases following inhibition of Hsp6 (mortalin/Grp75/mtHsp70) using RNAi.<sup>2</sup>

### References

1. Stiban, J. and Perera, M. Very long chain ceramides interfere with C16-ceramide-induced channel formation: A plausible mechanism for regulating the initiation of intrinsic apoptosis. *Biochim. Biophys. Acta* **1848**(2), 561-567 (2015).
2. Kim, H.E., Grant, A.R., Simic, M.S., et al. Lipid biosynthesis coordinates a mitochondrial-to-cytosolic stress response. *Cell* **166**(6), 1539-1552 (2016).

**WARNING**  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

**SAFETY DATA**  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

**WARRANTY AND LIMITATION OF REMEDY**  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/26/2021

### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM