

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Anacardic Acid diene

Item No. 22663

CAS Registry No.: 103904-74-1

Formal Name: 2-hydroxy-6-8Z,11Z-

pentadecadien-1-yl-benzoic acid

Synonym: Anacardic Acid 15:2

MF: $C_{22}H_{32}O_3$ FW: 344.5 ≥95% **Purity:**

UV/Vis.: λ_{max} : 244, 313 nm

Supplied as: A neat oil Storage: -20°C Stability: ≥1 year

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Anacardic acid diene is supplied as a neat oil. A stock solution may be made by dissolving the anacardic acid diene in the solvent of choice. Anacardic acid diene is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of anacardic acid diene in these solvents is approximately 22, 15, and 20 mg/ml, respectively.

Anacardic acid diene is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, anacardic acid diene should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Anacardic acid diene has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

Anacardic acid diene is a polyunsaturated form of anacardic acid (Item No. 13144) that has been found in cashew nut shell liquid. It has antibacterial activity against methicillin resistant S. aureus (MRSA) and S. mutans (MICs = 12.5 and 6.25 μg/ml, respectively).² Anacardic acid diene has schistosomicidal activity against adult S. mansoni worms when used at a concentration of 100 μM.¹ It also inhibits soybean lipoxygenase-1 in a time-dependent manner.3

References

- 1. Alvarenga, T.A., de Oliveira, P.F., de Souza, J.M., et al. Schistosomicidal activity of alkyl-phenols from the cashew Anacardium occidentale against Schistosoma mansoni adult worms. J. Agric. Food Chem. 64(46), 8821-8827 (2016).
- 2. Kubo, I., Nihei, K., and Tsujimoto, K. Antibacterial action of anacardic acids against methicillin resistant Staphylococcus aureus (MRSA). J. Agric. Food Chem. 51(26), 7624-7628 (2003).
- Ha, T.J. and Kubo, I. Lipoxygenase inhibitory activity of anacardic acids. J. Agric. Food Chem. 53(11), 4350-4354 (2005).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 09/12/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM