



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## Eicosapentaenoyl Chloride

Item No. 22721

**CAS Registry No.:** 98770-65-1  
**Formal Name:** (5Z,8Z,11Z,14Z,17Z)-5,8,11,14,17-eicosapentaenoyl chloride  
**Synonyms:** Eicosapentaenoic Acid Chloride, EPA Chloride  
**MF:** C<sub>20</sub>H<sub>29</sub>ClO  
**FW:** 320.9  
**Purity:** ≥95%  
**Supplied as:** A liquid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Eicosapentaenoyl chloride is supplied as a liquid. A stock solution may be made by dissolving the eicosapentaenoyl chloride in the solvent of choice, which should be purged with an inert gas. Eicosapentaenoyl chloride is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of eicosapentaenoyl chloride in these solvents is approximately 10 mg/ml.

### Description

Eicosapentaenoyl chloride is a derivative of eicosapentaenoic acid (Item Nos. 90110 | 90110.1 | 21908). It has been used in the synthesis of fatty acid conjugates to enhance lipophilicity and cell permeability of bioactive compounds such as (-)-epigallocatechin gallate (EGCG; Item No. 70935) and salicylic acid.<sup>1,2</sup>

### References

1. Zhong, Y., Chiou, Y.-S., Pan, M.-H., *et al.* Anti-inflammatory activity of lipophilic epigallocatechin gallate (EGCG) derivatives in LPS-stimulated murine macrophages. *Food Chem.* **134**(2), 742-748 (2012).
2. Vu, C.-B., Bemis, J.E., Benson, E., *et al.* Synthesis and characterization of fatty acid conjugates of niacin and salicylic acid. *J. Med. Chem.* **59**(3), 1217-1231 (2016).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/06/2020

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM