



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



C2 Phytoceramide (t18:0/2:0)

Item No. 22823

Formal Name: N-[(1S,2S,3R)-2,3-dihydroxy-1-(hydroxymethyl)heptadecyl]-acetamide

Synonyms: N-Acetyl Phytosphingosine, C2:0 Phytoceramide, Cer(t18:0/2:0), Ceramide (t18:0/2:0), NAPS

MF: C₂₀H₄₁NO₄


FW: 359.6

Purity: ≥98%

Supplied as: A solid

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

C2 Phytoceramide (t18:0/2:0) is supplied as a solid. A stock solution may be made by dissolving the C2 phytoceramide (t18:0/2:0) in the solvent of choice. C2 Phytoceramide (t18:0/2:0) is soluble in organic solvents such as ethanol, methanol, and DMSO.

Description

C2 Phytoceramide is a bioactive semisynthetic sphingolipid that inhibits formyl peptide-induced oxidant release (IC₅₀ = 0.38 μM) in suspended polymorphonuclear cells.¹ It increases COX-2 protein levels 15-fold through ERK signaling.² It induces death of keratinocytes (20% viability) with an ED₅₀ value of 30 μM, the same concentration at which 35% of cells in a TUNEL assay are apoptotic. C2 Phytoceramide also has antiproliferative effects in CHO cells, with greater than 80% cytotoxicity achieved at a concentration of 20 μM, and induces internucleosomal DNA fragmentation.³ In addition, it inhibits the activation of phospholipase D (PLD) mediated by muscarinic acetylcholine receptors *in vitro*.

References

1. Nakamura, T., Abe, A., Balazovich, K.J., *et al.* Ceramide regulates oxidant release in adherent human neutrophils. *J. Biol. Chem.* **269**(28), 18384-18389 (1994).
2. Kim, H.J., Shin, W., Park, C.S., *et al.* Differential regulation of cyclooxygenase-2 expression by phytosphingosine derivatives, NAPS and TAPS, and its role in the NAPS or TAPS-mediated apoptosis. *J. Invest. Dermatol.* **121**(5), 1126-1134 (2003).
3. Lee, J.S., Min, D.S., Park, C., *et al.* Phytosphingosine and C2-phytoceramide induce cell death and inhibit carbachol-stimulated phospholipase D activation in Chinese hamster ovary cells expressing the *Caenorhabditis elegans* muscarinic acetylcholine receptor. *FEBS. Lett.* **499**(1-2), 82-86 (2001).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/20/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM