

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



### Motolimod

Item No. 22952

CAS Registry No.: 926927-61-9

Formal Name: 2-amino-N,N-dipropyl-8-[4-(1-

pyrrolidinylcarbonyl)phenyl]-3H-


1-benzazepine-4-carboxamide

Synonym: VTX-2337 MF:  $C_{28}H_{34}N_4O_2$ FW: 458.6 **Purity:** ≥98%

UV/Vis.:  $\lambda_{max}$ : 270 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



#### **Laboratory Procedures**

Motolimod is supplied as a crystalline solid. A stock solution may be made by dissolving the motolimod in the solvent of choice. Motolimod is soluble in the organic solvent DMSO, which should be purged with an inert gas, at a concentration of approximately 2 mg/ml.

Motolimod is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, motolimod should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. Motolimod has a solubility of approximately 0.2 mg/ml in a 1:4 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

Motolimod is an agonist of toll-like receptor 8 (TLR8). It increases the production of TNF- $\alpha$  and IL-12 in human peripheral blood mononuclear cells (PBMCs;  $EC_{50}s = 140$  and 120 nM, respectively), monocytes, and myeloid dendritic cells. Motolimod also increases IFN-γ production in natural killer (NK) cells and increases cytolysis in K562 NK cell-sensitive leukemia cells when used at concentrations of 167 and 500 nM for 48 hours. It increases plasma levels of a group of human cytokines, including IL-6, IL-12p70, TNF-α, MCP-1, and MIP-1β, in NOD-scid IL2ry<sup>null</sup> (NSG) mice reconstituted with human immune system (NSG-HIS) when administered at doses of 1.5 and 15 mg/m<sup>2.2</sup> Motolimod, when used in combination with pegylated liposomal doxorubicin (PLD), reduces tumor growth and increases tumor infiltration of monocytes and T cells in an ovarian cancer NSG-HIS mouse model.

#### References

- 1. Lu, H., Dietsch, G.N., Matthews, M.A., et al. VTX-2337 is a novel TLR8 agonist that activates NK cells and augments ADCC. Clin Cancer Res. 18(2), 499-509 (2012).
- 2. Monk, B.J., Facciabene, A., Brady, W.E., et al. Integrative development of a TLR8 agonist for ovarian cancer chemoimmunotherapy. Clin Cancer Res. 23(8), 1955-1966 (2017).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

al should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution

## WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/19/2018

#### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM