

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



Cardanol monoene

Item No. 23154

CAS Registry No.: Formal Name: Synonym: MF: FW: Purity: UV/Vis.:	3-8Z-pentadecen-1-yl-phenol Cardanol C15:1 C <sub>21</sub> H <sub>34</sub> O 302.5 ≥98% λ <sub>max</sub> : 276 nm	
Supplied as:	A neat oil	
Storage:	-20°C	
Stability:	≥1 year	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

#### Laboratory Procedures

Cardanol monoene is supplied as a neat oil. A stock solution may be made by dissolving the oil in the solvent of choice. Cardanol monoene is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of cardanol monoene in these solvents is approximately 22, 15, and 20 mg/ml, respectively.

Cardanol monoene is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, cardanol monoene should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. Cardanol monoene has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

Cardanol monoene is a phenol found in cashew nut shell liquid that reversibly inhibits tyrosinase with an IC<sub>50</sub> value of 56  $\mu$ M in vitro.<sup>1</sup> It halts the cell cycle at the S phase and inhibits proliferation dose- and time-dependently in M14 melanoma cells (IC  $_{50}$ s = 23.15 and 12.30  $\mu$ M after 24 and 48 hours of treatment, respectively).<sup>2</sup> Cardanol monoene also induces apoptosis via the intrinsic pathway following the accumulation of reactive oxygen species (ROS). A mixture of cardanol mono-, di-, and triene is used to synthesize cardanolmetal complexes that inhibit uropathogenic E. coli biofilm formation.<sup>3</sup>

#### References

- 1. Yu, X.-P., Su, W.-C., Wang, Q., et al. Inhibitory mechanism of cardanols on tyrosinase. Proc. Biochem. 51(12), 2330-2237 (2016).
- 2. Su, W.-C., Lin, Y.-F., Wang, Y.-X., et al. Mitochondria-associated apoptosis in human melanoma cells induced by cardanol monoene from cashew nut shell liquid. J. Agric. Food Chem. 65(28), 5620-5631 (2017).
- 3. Lalitha, K., Prasad, Y.S., Sridharan, V., et al. Intrinsic hydrophobic antibacterial thin film from renewable resources: Application in the development of anti-biofilm urinary catheters. ACS Sus. Chem. Eng. (Univ. of Berlin) 5(1), 436 (2017).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/20/2017

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM