

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



### Tetrachlorohydroquinone

Item No. 23231

CAS Registry No.:	87-87-6	
Formal Name:	2,3,5,6-tetrachloro-1,4-benzenediol	ÇI
Synonyms:	NSC 4858, NSC 100888, TCHQ	
MF:	C <sub>6</sub> H <sub>2</sub> Cl <sub>4</sub> O <sub>2</sub>	HO
FW:	247.9	
Purity:	≥98%	
UV/Vis.:	λ <sub>max</sub> : 309 nm	СІ ОН
Supplied as:	A crystalline solid	
Storage:	-20°C	CI
Stability:	≥4 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

#### Laboratory Procedures

Tetrachlorohydroquinone (TCHQ) is supplied as a crystalline solid. A stock solution may be made by dissolving the TCHQ in the solvent of choice, which should be purged with an inert gas. TCHQ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of TCHQ in these solvents is approximately 30 mg/ml.

TCHQ is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, TCHQ should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. TCHQ has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

#### Description

TCHQ is a metabolite of the organochlorine biocide pentachlorophenol.<sup>1</sup> It is cytotoxic to RTL-W1 rainbow trout liver cells (EC<sub>50</sub> = 1.55  $\mu$ M in a neutral red assay). TCHQ increases production of reactive oxygen species (ROS), inhibits apoptosis, and induces loss of the mitochondrial membrane potential and necrosis in splenocytes.<sup>2</sup> In vivo, TCHQ induces glutathione (GSH) depletion in mouse liver.<sup>3</sup>

#### References

- 1. Pietsch, C., Hollender, J., Dorusch, F., et al. Cytotoxic effects of pentachlorophenol (PCP) and its metabolite tetrachlorohydroquinone (TCHQ) on liver cells are modulated by antioxidants. Cell Biol. Toxicol. 30(4), 233-252 (2014).
- 2. Chen, H.-M., Zhu, B.-Z., Chen, R.-J., et al. The pentachlorophenol metabolite tetrachlorohydroquinone induces massive ROS and prolonged p-ERK expression in splenocytes, leading to inhibition of apoptosis and necrotic cell death. PLoS One 9(2), e89483 (2014).
- 3. Wang, Y.-J., Ho, Y.-S., Chu, S.-W., et al. Induction of glutathione depletion, p53 protein accumulation and cellular transformation by tetrachlorohydroquinone, a toxic metabolite of pentachlorophenol. Chem. Biol. Interact. 105(1), 1-16 (1997).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/11/2022

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM