



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



β -Cyclodextrin (hydrate)

Item No. 23387

CAS Registry No.: 68168-23-0
Formal Name: β -cyclodextrin, hydrate
Synonyms: β -CD, NSC 269471, NSC 314334, Schardinger β -Dextrin
MF: $C_{42}H_{70}O_{35} \cdot XH_2O$
FW: 1,135.0
Purity: $\geq 95\%$
Supplied as: A crystalline solid
Storage: $-20^\circ C$
Stability: ≥ 2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

β -Cyclodextrin (β -CD) (hydrate) is supplied as a crystalline solid. A stock solution may be made by dissolving the β -CD (hydrate) in the solvent of choice. β -CD (hydrate) is soluble in organic solvents such as DMSO and dimethyl formamide, which should be purged with an inert gas. The solubility of β -CD (hydrate) in these solvents is approximately 2.5 mg/ml.

β -CD (hydrate) is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, β -CD (hydrate) should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. β -CD (hydrate) has a solubility of approximately 0.03 mg/ml in a 1:30 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

β -CD is a cyclic oligosaccharide that contains seven D-(+)-glucopyranose units and has been used to improve the aqueous solubility of various compounds, especially those containing a phenyl group.^{1,2} The circular arrangement of its glucose units produces a torus-shaped ring configuration in which the CH_2 groups and ether linkages of the molecule face the hollow interior, resulting in a nonpolar, hydrophobic cavity and a polar, hydrophilic exterior. When combined in solution with other compounds, the nonpolar aromatic portions of that compound interact with the nonpolar interior of the β -CD molecule, thus isolating the aromatic portion of the molecule from the water and thereby increasing its aqueous solubility.^{1,3}

References

1. Katageri, A.R. and Sheikh, M.A. Cyclodextrin a gift to pharmaceutical world review. *Int. Res. J. Pharm.* **3(1)**, 52-56 (2012).
2. Williamson, G.B. and Weidenhamer, J.D. Bacterial degradation of juglone: Evidence against allelopathy? *J. Chem. Ecol.* **16(5)**, 1739-1742 (1990).
3. Rao, V.M., Nerurkar, M., Pinnamaneni, S., et al. Co-solubilization of poorly soluble drugs by micellization and complexation. *Int. J. Pharm.* **319(1-2)**, 98-106 (2006).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/29/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM