



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION




Reveromycin C

Item No. 23488

CAS Registry No.: 144860-69-5
Formal Name: butanedioic acid, 1-[(2S,3R,6S,8R,9S)-8-[(2E,4E,6S,7S,8E)-9-carboxy-6-hydroxy-3,7-dimethyl-2,4,8-nonatrien-1-yl]-2-[(1E,3E)-4-carboxy-3-methyl-1,3-butadien-1-yl]-9-methyl-3-(3-methylbutyl)-1,7-dioxaspiro[5.5]undec-3-yl] ester

MF: C₃₇H₅₄O₁₁
FW: 674.8
Purity: ≥95%
Supplied as: A lyophilisate
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Reveromycin C is supplied as a lyophilisate. A stock solution may be made by dissolving the reveromycin C in the solvent of choice. Reveromycin C is soluble in organic solvents such as ethanol, methanol, DMSO, and dimethyl formamide, which should be purged with an inert gas.

Reveromycin C is sparingly soluble in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

Description

Reveromycin C is a polyketide originally isolated from *Streptomyces* that has antifungal activity against *C. albicans* (MICs = 2.0 and >500 µg/ml at pH 3 and 7.4, respectively).^{1,2} Reveromycin C inhibits EGF-induced mitogenic activity in the BALB/MK mouse epidermal cell line.¹ It also reverses the morphology of sarcoma-virus-transformed NRK rat kidney cells (EC₅₀ = 1.58 µg/ml) and inhibits proliferation of KB cells and K562 human chronic myelogenous leukemia cells (IC₅₀ = 2.0 µg/ml for both).¹

References

1. Miyazawa, T., Takahashi, S., Kawata, A., *et al.* Identification of middle chain fatty Acyl-CoA ligase responsible for the biosynthesis of 2-alkylmalonyl-CoAs for polyketide extender unit. *J. Biol. Chem.* **290**(45), 26994-27011 (2015).
2. Takahashi, H., Osada, H., Koshino, H., *et al.* Reveromycins, new inhibitors of eukaryotic cell growth. II. Biological activities. *J. Antibiot. (Tokyo)* **45**(9), 1414-1419 (1992).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/25/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM