



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## Dipivefrin (hydrochloride)

Item No. 24028

**CAS Registry No.:** 64019-93-8  
**Formal Name:** 2,2-dimethyl-propanoic acid, 1,1'-[4-[1-hydroxy-2-(methylamino)ethyl]-1,2-phenylene] ester, monohydrochloride  
**MF:** C<sub>19</sub>H<sub>29</sub>NO<sub>5</sub> • HCl  
**FW:** 387.9  
**Purity:** ≥98%  
**Supplied as:** A crystalline solid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

Dipivefrin (hydrochloride) is supplied as a crystalline solid. A stock solution may be made by dissolving the dipivefrin (hydrochloride) in the solvent of choice. Dipivefrin (hydrochloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of dipivefrin (hydrochloride) in these solvents is approximately 33 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of dipivefrin (hydrochloride) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of dipivefrin (hydrochloride) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

Dipivefrin is a prodrug of epinephrine that is hydrolyzed by cholinesterase and other esterases in the cornea to epinephrine.<sup>1</sup> It reduces the density of cultured bovine primary trabecular meshwork cells (IC<sub>50</sub> = 115 μM).<sup>2</sup> Additionally, it induces an elongated, fibroblast-like morphology and disrupts the actin cytoskeleton in bovine primary trabecular meshwork cells when used at a concentration of 103 μM.<sup>2</sup> In cultured bovine corneal endothelial cells, dipivefrin (28 μM) enhances calcium signaling and induces cytotoxicity.<sup>3,4</sup> Dipivefrin also suppresses primary human corneal keratinocyte proliferation when used at a concentration of 280 μM.<sup>5</sup> Formulations containing dipivefrin have been used alone and in combination with β-adrenergic receptor antagonists for the treatment of glaucoma.

### References

1. Nakamura, M., Shirasawa, E., and Hikida, M. Characterization of esterases involved in the hydrolysis of dipivefrin hydrochloride. *Ophthalmic Res.* **25(1)**, 46-51 (1993).
2. Kawa, J.E., Higginbotham, E.J., Change, I.L., *et al.* Effects of antiglaucoma medications on bovine trabecular meshwork cells in vitro. *Exp. Eye Res.* **57(5)**, 557-565 (1993).
3. Wu, K.Y., Hong, S.J., and Wang, H.Z. Effects of antiglaucoma drugs on calcium mobility in cultured corneal endothelial cells. *Kaohsiung J. Med. Sci.* **22(2)**, 60-67 (2006).
4. Wu, K.Y., Wang, H.Z., and Hong, S.J. Cellular cytotoxicity of antiglaucoma drugs in cultured corneal endothelial cells. *Kaohsiung J. Med. Sci.* **23(3)**, 105-111 (2007).
5. Wu, K.Y., Wang, H.Z., and Hong, S.J. Effects of antiglaucoma drugs on cellular proliferation in cultured human corneal keratocytes. *Kaohsiung J. Med. Sci.* **22(3)**, 120-125 (2006).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/15/2018

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM