



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



C8 dihydro Ceramide (d18:0/8:0)

Item No. 24358

CAS Registry No.: 145774-33-0
Formal Name: N-[(1S,2R)-2-hydroxy-1-(hydroxymethyl)heptadecyl]-octanamide
Synonyms: Cer(d18:0/8:0), Ceramide (d18:0/8:0), N-octanoyl-D-erythro-Dihydrosphingosine
MF: C₂₆H₅₃NO₃
FW: 427.7
Purity: ≥98%
Supplied as: A solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

C8 dihydro Ceramide (d18:0/8:0) is supplied as a solid. A stock solution may be made by dissolving the C8 dihydro ceramide (d18:0/8:0) in the solvent of choice. C8 dihydro Ceramide (d18:0/8:0) is soluble in organic solvents such as ethanol, DMSO, and chloroform, which should be purged with an inert gas.

Description

C8 dihydro Ceramide is a sphingolipid and an intermediate in the synthesis of C8 ceramide (Item No. 62540).¹ It is synthesized by the acylation of sphinganine by ceramide synthase, a process that can be inhibited by some fungal mycotoxins, such as fumonisin B₁ (Item No. 62580).² C8 dihydro Ceramide can be converted to C8 ceramide *via* the introduction of a 4,5-*trans* double bond by dihydroceramide desaturase.¹ C8 dihydro Ceramide is metabolically inactive and has been used as a negative control for the biological activity of C8 ceramide.³

References

1. Michel, C., van Echten-Deckert, G., Rother, J., *et al.* Characterization of ceramide synthesis. A dihydroceramide desaturase introduces the 4,5-*trans*-double bond of sphingosine at the level of dihydroceramide. *J. Biol. Chem.* **272(36)**, 22432-22437 (1997).
2. Wang, E., Norred, W.P., Bacon, C.W., *et al.* Inhibition of sphingolipid biosynthesis by fumonisins. Implications for diseases associated with *Fusarium moniliforme*. *J. Biol. Chem.* **266(22)**, 14486-14490 (1991).
3. Struckhoff, A.P., Bittman, R., Burow, M.E., *et al.* Novel ceramide analogs as potential chemotherapeutic agents in breast cancer. *J. Pharmacol. Exp. Ther.* **309(2)**, 523-532 (2004).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/14/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM