



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



L-erythro Sphinganine (d18:0)

Item No. 24374

CAS Registry No.: 6036-76-6
Formal Name: 2R-amino-1,3S-octadecanediol
Synonyms: L-erythro Dihydrosphingosine,
C18 L-erythro Sphinganine (d18:0)
MF: C₁₈H₃₉NO₂
FW: 301.5
Purity: ≥98%
Supplied as: A solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

L-erythro Sphinganine (d18:0) is supplied as a solid. A stock solution may be made by dissolving the L-erythro sphinganine (d18:0) in the solvent of choice. L-erythro Sphinganine (d18:0) is soluble in organic solvents such as chloroform, methanol, ethanol, and DMSO, which should be purged with an inert gas.

Description

L-erythro Sphinganine (d18:0) is a synthetic bioactive sphingolipid and stereoisomer of sphinganine (d18:0) (Item No. 10007945) and D-threo sphinganine (d18:0) (Item No. 24375).¹ It induces autophagy in HCT116 cells when used at a concentration of 12 μM. L-erythro Sphinganine induces cell cycle arrest of 7R4-LCB1 yeast cells.¹ It is metabolized via sphinganine N-acyltransferase and sphinganine kinase *in vivo* in rat liver.²

References

1. Coward, J., Ambrosini, G., Musi, E., *et al.* Safingol (L-threo-sphinganine) induces autophagy in solid tumor cells through inhibition of PKC and the PI3-kinase pathway. *Autophagy* 5(2), 184-193 (2009).
2. Stoffel, W. and Bister, K. Stereospecificities in the metabolic reactions of the four isomeric sphingamines (dihydrosphingosines) in rat liver. *Hoppe-Seylers Z. Physiol. Chem.* 354(2), 169-181 (1973).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/13/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM