

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



C16 Glucosylceramide-d₃ (d18:1/16:0-d₃)

Item No. 24621

Formal Name: N-((2S,3R,E)-3-hydroxy-1-

> (((2R,3R,4S,5S,6R)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2Hpyran-2-yl)oxy)octadec-4-en-2-yl)

hexadecanamide-d₃

Synonyms: GluCer(d18:1/16:0-d₃),

Glucosylceramide-d₃ (d18:1/16:0-d₃)

 $C_{40}H_{74}D_3NO_8$ MF:

FW: 703.1


Chemical Purity: ≥98% (C16 Glucosylceramide)

Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₃); \leq 1% d₀

Supplied as: A solid -20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

C16 Glucosylceramide-d₃ (d18:1/16:0-d₃) is supplied as a solid. A stock solution may be made by dissolving the C16 glucosylceramide-d₃ (d18:1/16:0-d₃) in the solvent of choice. C16 Glucosylceramide-d₃ (d18:1/16:0-d₃) is soluble in a 2:1 solution of chloroform:methanol.

Description

C16 Glucosylceramide-d3 is intended for use as an internal standard for the quantification of C16 glucosylceramide by GC- or LC-MS. C16 glucosylceramide is an endogenous bioactive sphingolipid. 1,2 It is a precursor in the synthesis of C16 lactosylceramide (Item No. 24352) that is formed via metabolism of C16 ceramide (Item No. 10681) by glucosylceramide synthase.³ Inhalation of C16 glucosylceramide reduces lung colonization by P. aeruginosa and increases survival in a mouse model of cystic fibrosis.1 C16 Glucosylceramide levels are elevated in the plasma of Parkinson's disease patients with cognitive impairments.2

References

- 1. Kovacic, B., Sehl, C., Wilker, B., et al. Glucosylceramide critically contributes to the host defense of cystic fibrosis lungs. Cell Physiol. Biochem. 41(3), 1208-1218 (2017).
- 2. Mielke, M.M., Maetzler, W., Haughey, N.J., et al. Plasma ceramide and glucosylceramide metabolism is altered in sporadic Parkinson's disease and associated with cognitive impairment: A pilot study. PLoS One 8(9), e73094 (2013).
- 3. Holland, W.L. and Summers, S.A. Sphingolipids, insulin resistance, and metabolic disease: New insights from in vivo manipulation of sphingolipid metabolism. Endocr. Rev. 29(4), 381-402 (2008).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 03/19/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM