

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# **PRODUCT INFORMATION**



## **ODG**

Item No. 24782

Formal Name: N-(2-methyl-1,3-bis((3R,4S,5S-

> trihydroxy-6R-(hydroxymethyl) tetrahydro-2H-pyran-2R-yl)

oxy)propan-2-yl)-3-(octylthio)

propanamide

MF: C<sub>27</sub>H<sub>51</sub>NO<sub>13</sub>S FW:

629.8 **Purity:** ≥95% Supplied as: A powder Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### **Laboratory Procedures**


ODG is supplied as a powder. A stock solution may be made by dissolving the ODG in the solvent of choice, which should be purged with an inert gas. ODG is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of ODG in ethanol and DMSO is approximately 10 mg/ml and . approximately 2 mg/ml in DMF

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of ODG can be prepared by directly dissolving the powder in aqueous buffers. The solubility of ODG in PBS, pH 7.2, is approximately 2 mg/ml. We do not recommend storing the aqueous solution for more than one day.

### Description

ODG is a detergent used to solubilize membrane proteins. It has a critical micelle concentration (CMC) of greater than 10 mM.

#### **Images**



Thermostability curves 1,2 Normalyzed intensity 0,8 (oC) 0,6 20 ODG 0.2 0,0 -0,2 Temperature (°C)

Stabilization of GPCR target. The GPCR protein was extracted using either reference detergent or ODG and heated at different temperatures for 30 min. After centrifugation at 20,000 g for 40 min, samples were separated on a 4-15% Tris-glycine SDS-PAGE, transferred to PVDF membrane and immunodetected with a specific antibody. Band intensity was measured and the resulting graph

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/12/2019

#### **CAYMAN CHEMICAL**

Tm

ODG

Reference

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM