



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



17-phenyl trinor Prostaglandin F_{2α} isopropyl amide

Item No. 25242

CAS Registry No.: 155205-99-5
Formal Name: (Z)-7-((1R,2R,3R,5S)-3,5-dihydroxy-2-((S,E)-3-hydroxy-5-phenylpent-1-en-1-yl)cyclopentyl)-N-isopropylhept-5-enamide
Synonym: 17-phenyl trinor PGF_{2α} isopropyl amide
MF: C₂₆H₃₉NO₄
FW: 429.6
Purity: ≥98%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

17-phenyl trinor Prostaglandin F_{2α} isopropyl amide (17-phenyl trinor PGF_{2α} isopropyl amide) is supplied as a crystalline solid. A stock solution may be made by dissolving the 17-phenyl trinor PGF_{2α} isopropyl amide in the solvent of choice, which should be purged with an inert gas. 17-phenyl trinor PGF_{2α} isopropyl amide is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of 17-phenyl trinor PGF_{2α} isopropyl amide in ethanol is approximately 50 mg/ml and approximately 25 mg/ml in DMSO and DMF.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 17-phenyl trinor PGF_{2α} isopropyl amide can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of 17-phenyl trinor PGF_{2α} isopropyl amide in PBS, pH 7.2, is approximately 0.3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

17-phenyl trinor PGF_{2α} isopropyl amide is a derivative of 17-phenyl trinor PGF_{2α} ethyl amide (Item No. 16820).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/24/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM