

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



#### **GSK2033**

Item No. 25443

CAS Registry No.: Formal Name:	1221277-90-2 2,4,6-trimethyl-N-[[3'- (methylsulfonyl)[1,1'-biphenyl]-4-	
	yl]methyl]-N-[[5-(trifluoromethyl)-	
	2-furanyl]methyl]-	
MF:	$C_{29}H_{28}F_3NO_5S_2$	
FW:	591.7	
Purity:	≥98%	
UV/Vis.:	λ <sub>max</sub> : 255 nm	S S
Supplied as:	A solid	
Storage:	-20°C	
Stability:	≥2 years	

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

#### Laboratory Procedures

GSK2033 is supplied as a solid. A stock solution may be made by dissolving the GSK2033 in the solvent of choice, which should be purged with an inert gas. GSK2033 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of GSK2033 in these solvents is approximately 3 mg/ml for DMSO and 5 mg/ml for DMF. GSK2033 is slightly soluble in ethanol.

#### Description

GSK2033 is an antagonist of liver X receptor  $\alpha$  (LXR $\alpha$ ) and LXR $\beta$  (IC<sub>50</sub>S = 0.1 and 0.398  $\mu$ M, respectively) that has no agonist activity in an LXR transactivation assay.<sup>1</sup> It inhibits LXR agonist-induced and basal expression of the LXR target genes ATP-binding cassette transporter 1 (ABCA1) and sterol regulatory element binding protein 1c (SREBP-1c) in THP-1 and HepG2 cells, respectively. However, GSK2033 binds promiscuously to a variety of nuclear receptors including ROR $\gamma$ , RXR $\alpha$ , ER $\alpha$ , and ER $\beta$  in a nuclear receptor specificity assay in HEK293 cells and, in a mouse model of non-alcoholic fatty liver disease (NAFLD), it induces the expression of fatty acid synthase and SREBP-1.<sup>2</sup>

#### References

- 1. Zuercher, W.J., Buckholz, R.G., Campobasso, N., et al. Discovery of tertiary sulfonamides as potent liver X receptor antagonists. J. Med Chem. 53(8), 3412-3416 (2010).
- 2. Griffett, K. and Burris, T.P. Promiscuous activity of the LXR antagonist GSK2033 in a mouse model of fatty liver disease. Biochem. Biophys. Res. Commun. 479(3), 424-428 (2016).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

SAFETY DATA This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/08/2021

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM