



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



C6 Urea Ceramide

Item No. 25491

CAS Registry No.: 486991-52-0
Formal Name: N-hexyl-N'-[(1S,2R,3E)-2-hydroxy-1-(hydroxymethyl)-3-heptadecen-1-yl]-urea
Synonyms: C6 Ceramide (d18:1/6:0) Urea, Cer(d18:1/6:0) Urea, D-erythro-Urea-C6-Ceramide
MF: C₂₅H₅₀N₂O₃
FW: 426.7
Purity: ≥95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

C6 urea ceramide is supplied as a crystalline solid. A stock solution may be made by dissolving the C6 urea ceramide in the solvent of choice, which should be purged with an inert gas. C6 urea ceramide is soluble in the organic solvent dimethyl formamide at a concentration of approximately 0.5 mg/ml.

Description

C6 urea ceramide is an inhibitor of neutral ceramidase.¹ It increases total ceramide levels in wild-type mouse embryonic fibroblasts (MEFs) and in HT-29 colon cancer cells but not in MEFs lacking neutral ceramidase. It inhibits proliferation of, and induces apoptosis and autophagy in HT-29, but not non-cancerous RIE-1, cells when used at concentrations of 5 and 10 μM. C6 urea ceramide decreases total β-catenin, increases phosphorylated β-catenin, and induces colocalization of β-catenin with the 20S proteasome in HT-29 and HCT116, but not RIE-1, cells. It reduces tumor growth and increases C16, C18, C20, and C24 ceramide levels in tumor tissue in an HT-29 mouse xenograft model when administered at doses of 1.25, 2.5, and 5 mg/kg for five days.

Reference

1. García-Barros, M., Coant, N., Kawamori, T., *et al.* Role of neutral ceramidase in colon cancer. *FASEB J.* **30**(12), 4159-4171 (2016).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/14/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM