

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



(R,S)-BHFF

Item No. 25535

CAS Registry No.: 123557-91-5

Formal Name: 5,7-bis(1,1-dimethylethyl)-3-hydroxy-3-

(trifluoromethyl)-2(3H)-benzofuranone

Synonym: MF: $C_{17}H_{21}F_3O_3$ 330.3 FW: ≥95% **Purity:**

UV/Vis.: λ_{max} : 285 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

(R,S)-BHFF is supplied as a crystalline solid. A stock solution may be made by dissolving the (R,S)-BHFF in the solvent of choice. (R,S)-BHFF is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of (R,S)-BHFF in these solvents is approximately 30 mg/ml.

(R,S)-BHFF is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, (R,S)-BHFF should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. (R,S)-BHFF has a solubility of approximately 0.25 mg/ml in a 1:3 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

(R,S)-BHFF is a positive allosteric modulator of GABA_B receptors that increases the potency of GABA by 15.3-fold in a GTP γ [35S] binding assay when used at a concentration of 0.3 μ M.¹ It is selective for GABA_B over a panel of receptors and ion channels at 10 μ M, however, it inhibits the cholecystokinin-1 (CCK_{Λ}) receptor (IC₅₀ = 4.1 μM). (R,S)-BHFF enhances inhibition of the population spike of CA1 pyramidal cells induced by the GABA_R receptor antagonist baclofen (Item No. 18600) in rat hippocampal slices. In vivo, (R,S)-BHFF increases the baclofen-induced loss of righting reflex and reverses stress-induced hyperthermia in mice. (R,S)-BHFF (50, 100, and 200 mg/kg) reduces the number of lever responses to alcohol by 30, 65, and 90%, respectively, and increases latency to the first response on the alcohol lever in Sardinian alcohol preferring (sP) rats when administered pre-conditioning at a dose of 200 mg/kg.²

References

- 1. Malherbe, P., Masciardi, R., Norcross, R.D., et al. Characterization of (R,S)-5,7-di-tert-butyl-3-hydroxy-3-trifluoromethyl-3H-benzofuran-2-one as a positive allosteric modulator of GABA_R receptors. Br. J. Pharmacol. 154(4), 797-811 (2008).
- 2. Maccioni, P., Thomas, A.W., Carai, M.A., et al. The positive allosteric modulator of the GABA_R receptor, rac-BHFF, suppresses alcohol self-administration. Drug Alcohol Depend. 109(1-3), 96-103 (2010).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 07/13/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM