

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



5α,6α-epoxy Cholestanol

Item No. 25539

CAS Registry No.: 1250-95-9

Formal Name: $(3\beta,5\alpha,6\alpha)$ -5,6-epoxy-cholestan-3-ol

Synonyms: Cholesterol 5a,6a-epoxide,


5α,6α-Epoxycholesterol, NSC 18176

MF: $C_{27}H_{46}O_{2}$ FW: 402.7 **Purity:** ≥95%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 vears

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

5α,6α-epoxy Cholestanol is supplied as a crystalline solid. A stock solution may be made by dissolving the 5α,6α-epoxy cholestanol in the solvent of choice. 5α,6α-epoxy Cholestanol is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of 5α , 6α -epoxy cholestanol in these solvents is approximately 20, 0.1, and 2 mg/ml, respectively.

5α,6α-epoxy Cholestanol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 5α,6α-epoxy cholestanol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 5α,6α-epoxy Cholestanol has a solubility of approximately 0.3 mg/ml in a 1:2 solution of ethanol: PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

5α,6α-epoxy Cholestanol is an oxysterol and a metabolite of cholesterol produced by oxidation.¹ 5α,6α-epoxy Cholestanol accumulates in MCF-7 breast cancer cells in a reactive oxygen species-dependent manner following tamoxifen and PBPE application and induces triacylglycerol biosynthesis by binding to liver X receptor β (LXR β). 2 5 α ,6 α -epoxy Cholestanol levels are increased in rat aorta and mesenteric artery following orchidectomy, an effect that can be prevented by a DHA-supplemented diet. Levels are also increased in post-mortem frontal and occipital cortex of patients with Alzheimer's disease.³

References

- 1. Villalpando, D.M., Rojas, M.M., García, H.S., et al. Dietary docosahexaenoic acid supplementation prevents the formation of cholesterol oxidation products in arteries from orchidectomized rats. PLoS One 12(10), e0185805 (2017).
- 2. Segala, G., de Medina, P., Iuliano, L., et al. 5,6-Epoxy-cholesterols contribute to the anticancer pharmacology of tamoxifen in breast cancer cells. Biochem. Pharmacol. 86(1), 175-189 (2013).
- Testa, G., Staurenghi, E., Zerbinati, C., et al. Changes in brain oxysterols at different stages of Alzheimer's disease: Their involvement in neuroinflammation. Redox Biol. 10, 24-33 (2016).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/12/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM