



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



1-Oleoyl-2-hydroxy-*sn*-glycero-3-PE

Item No. 25596

CAS Registry No.: 89576-29-4
Formal Name: 1-oleoyl-2-hydroxy-*sn*-glycero-3-phosphatidylethanolamine
Synonyms: 18:1 LPE, 18:1 Lyso-PE, 1-Oleoyl-2-hydroxy-*sn*-glycero-3-phosphoethanolamine
MF: C₂₃H₄₆NO₇P
FW: 479.6
Purity: ≥95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

1-Oleoyl-2-hydroxy-*sn*-glycero-3-PE is supplied as a crystalline solid. A stock solution may be made by dissolving the 1-oleoyl-2-hydroxy-*sn*-glycero-3-PE in the solvent of choice. 1-Oleoyl-2-hydroxy-*sn*-glycero-3-PE is soluble in the organic solvent chloroform, which should be purged with an inert gas, at a concentration of approximately 3 mg/ml.

Description

1-Oleoyl-2-hydroxy-*sn*-glycero-3-PE is a naturally-occurring lysophospholipid and an analog of plasmalogen lysophosphatidylethanolamine.¹⁻³ It induces transient increases in intracellular calcium in PC12 rat neuronal cells in a concentration-dependent manner, an effect that can be blocked by the lysophosphatidic acid receptor 1 (LPA₁) antagonist AM095 (Item No. 22141).² 1-Oleoyl-2-hydroxy-*sn*-glycero-3-PE (100 ng/ml) increases IL-2 production in CD1d-stimulated murine natural killer T (NKT) cell hybridoma 2H4 cells.³ It also induces production of IL-4, but not IFN- γ , in murine splenocytes when used at a concentration of 100 ng/ml and in mouse serum when administered at a dose of 1 μ g per animal, indicating induction of the T helper 2 (Th2) response. 1-Oleoyl-2-hydroxy-*sn*-glycero-3-PE levels increase in interscapular brown adipose tissue (iBAT) in mice following three-day cold exposure.¹

References

1. Marcher, A.B., Loft, A., Nielsen, R., *et al.* RNA-Seq and mass-spectrometry-based lipidomics reveal extensive changes of glycerolipid pathways in brown adipose tissue in response to cold. *Cell. Rep.* **13(9)**, 2000-2013 (2015).
2. Lee, J.M., Park, S.J., and Im, D.S. Lysophosphatidylethanolamine increases intracellular Ca²⁺ through LPA₁ in PC-12 neuronal cells. *Biochem. Bioph. Res. Commun.* **461(2)**, 378-382 (2015).
3. Ni, G., Li, Z., Liang, K., *et al.* Synthesis and evaluation of immunostimulant plasmalogen lysophosphatidylethanolamine and analogues for natural killer T cells. *Bioorg. Med. Chem.* **22(11)**, 2966-2973 (2014).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/04/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM