

## Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# **PRODUCT** INFORMATION



#### Cholestenone

Item No. 25713

CAS Registry No.:	601-57-0	
Formal Name:	cholest-4-en-3-one	$\mathbf{\hat{x}}$
Synonyms:	4-Cholesten-3-one, $\Delta^4$ -Cholestenone,	`
	NSC 63000, NSC 134926	
MF:	C <sub>27</sub> H <sub>44</sub> O	
FW:	384.6	
Purity:	≥95%	
UV/Vis.:	λ <sub>max</sub> : 242 nm	Ĥ Ĥ
Supplied as:	A crystalline solid	
Storage:	-20°C	0, ~ ~
Stability:	≥2 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.		

#### Laboratory Procedures

Cholestenone is supplied as a crystalline solid. A stock solution may be made by dissolving the cholestenone in the solvent of choice. Cholestenone is soluble in organic solvents such as ethanol and dimethyl formamide, which should be purged with an inert gas. The solubility of cholestenone in these solvents is approximately 2 and 0.1 mg/ml, respectively.

#### Description

Cholestanone is a cholesterol metabolite that has a keto group in place of the 3-hydroxy group on cholesterol.<sup>1</sup> It decreases TGF-β-induced Smad2 phosphorylation and TGF-β expression and prevents inhibition of DNA synthesis by TGF- $\beta$  in Mv1Lu cells when used at a concentration of 50  $\mu$ g/ml.<sup>2</sup> Increased fecal excretion of cholestenone is correlated with an increased risk of colorectal cancer. Cholestenone reduces serum cholesterol levels in a variety of animal models but is toxic to rats when administered at doses of 700-1,000 mg/kg per day, inducing hypertrophy in and reducing the activity of the adrenal gland.<sup>3</sup> It has been used as synthetic intermediate in the synthesis of steroids.<sup>4</sup>

#### References

- 1. Salen, G., Batta, A.K., Tint, G.S., et al. Inverse relationship between plasma cholestanol concentrations and bile acid synthesis in sitosterolemia. J. Lipid Res. 35(10), 1878-1887 (1994).
- 2. Chen, C.-L., Wu, D.-C., Liu, M.-Y., et al. Cholest-4-en-3-one attenuates TGF-β responsiveness by inducing TGF- $\beta$  receptors degradation in Mv1Lu cells and colorectal adenocarcinoma cells. J. Recept. Signal Transduct. Res. 37(2), 189-199 (2017).
- 3. Steinberg, D., Fredrickson, D.S., and Avigan, J. Effects of  $\Delta^4$ -cholestenone in animals and in man. Proc. Soc. Exp. Biol. Med. 97(4), 784-790 (1958).
- 4. Wu, K., Li, W., Song, J., et al. Production, purification, and identification of cholest-4-en-3-one produced by cholesterol oxidase from Rhodococcus sp. in aqueous/organic biphasic system. Biochem. Insights 8(Suppl 1), 1-8 (2015).

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 01/22/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM