



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PRODUCT INFORMATION



proTAME

Item No. 25835

CAS Registry No.: 1362911-19-0
Formal Name: (2S)-2-[[[4-methylphenyl)sulfonyl]amino]-9,13-dioxo-14-phenyl-7-[[[(2-phenylacetyl)oxy]methoxy]carbonyl]amino]-10,12-dioxo-6,8-diazatetradec-6-enoic acid, methyl ester

Synonym: Pro-N-4-tosyl-L-arginine methyl ester

MF: C₃₄H₃₈N₄O₁₂S


FW: 726.8

Purity: ≥90%

Supplied as: A solution in DMSO

Storage: -20°C

Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Description

proTAME is a cell-permeable prodrug form of N-4-tosyl-L-arginine methyl ester (TAME; Item No. 17550), an inhibitor of the anaphase-promoting complex/cyclosome (APC/C), that is converted to TAME by intracellular esterases.^{1,2} proTAME (12 μM) blocks association of APC/C with the activator CDH1 and inhibits degradation of APC/C substrates in HeLa cells.¹ It induces mitotic arrest in metaphase followed by cell death in synchronized HeLa H2B-GFP cells when used at a concentration of 12 μM. It also increases mitotic duration in asynchronous HeLa H2B-GFP cells at a concentration of 4 μM, an effect that is enhanced by knockdown of the APC/C co-activator CDC20. proTAME decreases the viability of several laboratory and primary patient-derived human multiple myeloma (MM) cell lines (IC₅₀s = 2.8-20.3 μM) and increases apoptosis in RPMI-8226, LP-1, NCI-H929, and U266 MM cells when used at a concentration of 12 μM.²

References

1. Zeng, X., Sigoillot, F., Gaur, S., *et al.* Pharmacologic inhibition of the anaphase-promoting complex induces a spindle checkpoint-dependent mitotic arrest in the absence of spindle damage. *Cancer Cell* **18(4)**, 382-395 (2010).
2. Lub, S., Maes, A., De Veirman, K., *et al.* Inhibiting the anaphase promoting complex/cyclosome induces a metaphase arrest and cell death in multiple myeloma cells. *Oncotarget* **7(4)**, 4062-4076 (2016).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 09/28/2022

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM