

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



1-Stearoyl-2-Arachidonoyl-sn-glycero-3-PE

Item No. 25871

CAS Registry No.: 61216-62-4

Formal Name: 5Z,8Z,11Z,14Z-eicosatetraenoic acid, (1R)-1-

1-Stearoyl-2-Arachidonoyl-sn-

[[[(2-aminoethoxy)hydroxyphosphinyl]oxy]

methyl]-2-[(1-oxooctadecyl)oxy]ethyl ester

Synonyms: 18:0/20:4-PE, PE(18:0/20:4), SAPE,

> glycero-3-Phosphatidylethanolamine, 1-Stearoyl-2-Arachidonoyl-sn-glycero-3-


Phosphoethanolamine

MF: $C_{43}H_{78}NO_8P$ 768.1 FW: **Purity:** ≥95%

Supplied as: A solution in methanol

-20°C Storage: Stability: ≥1 year Special Conditions: Light sensitive

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

1-Stearoyl-2-arachidonoyl-sn-glycero-3-PE is supplied as a solution in methanol. To change the solvent, simply evaporate the chloroform under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as ethanol purged with an inert gas can be used. The solubility of 1-stearoyl-2-arachidonoyl-sn-glycero-3-PE in ethanol is approximately 30 mg/ml.

1-Stearoyl-2-arachidonoyl-sn-glycero-3-PE is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, the ethanolic solution of 1-stearoyl-2-arachidonoyl-sn-glycero-3-PE should be diluted with the aqueous buffer of choice. 1-Stearoyl-2-arachidonoyl-sn-glycero-3-PE has a solubility of approximately 0.3 mg/ml in a 1:2 solution of ethanol:PBS (pH 7.2) using this method.

Description

1-Stearoyl-2-arachidonoyl-sn-glycero-3-PE is a naturally-occurring phospholipid that is formed via the phosphatidylserine decarboxylation pathway in mammalian cells. It has been used in the generation of giant unilamellar vesicles for use in the study of biological membranes.²

References

- 1. Bleijerveld, O.B., Brouwers, J.F., Vaandrager, A.B., et al. The CDP-ethanolamine pathway and phosphatidylserine decarboxylation generate different phosphatidylethanolamine molecular species. J. Biol. Chem. 282(39), 28362-28372 (2007).
- 2. Billerit, C., Jeffries, G.D.M., Orwar, O., et al. Formation of giant unilamellar vesicles from spin-coated lipid films by localized IR heating. Soft Matter 8(42), 10823-10826 (2012).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/20/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM