



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



5-OxoETE methyl ester

Item No. 26059

CAS Registry No.: 74785-00-5
Formal Name: 5-oxo-6E,8Z,11Z,14Z-eicosatetraenoic acid, methyl ester
Synonym: 5-KETE methyl ester
MF: C₂₁H₃₂O₃
FW: 332.5
Purity: ≥95%
UV/Vis.: λ_{max}: 280 nm
Supplied as: A solution in ethanol
Storage: -80°C
Stability: ≥1 year



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

5-OxoETE methyl ester is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. 5-OxoETE methyl ester is miscible in these solvents.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. If an organic solvent-free solution of 5-oxoETE methyl ester is needed, it can be prepared by evaporating the ethanol and directly dissolving the neat oil in aqueous buffers. The solubility of 5-oxoETE methyl ester in PBS, pH 7.2, is approximately 0.8 mg/ml. For greater aqueous solubility, 5-oxoETE methyl ester can be directly dissolved in 0.1 M Na₂CO₃ (2 mg/ml) and then diluted with PBS (pH 7.2) to achieve the desired concentration or pH. We do not recommend storing the aqueous solution for more than one day.

Description

5-OxoETE methyl ester is an esterified form of the polyunsaturated keto acid 5-oxoETE (Item No. 34250). 5-OxoETE methyl ester is an agonist of oxoeicosanoid receptor 1 (OXER1; EC₅₀ = 1.54 μM for β-arrestin recruitment) that has a higher maximal response than 5-oxoETE in a β-arrestin assay.¹

Reference

- Stepniewski, T.M., Torrens-Fontanals, M., Rodríguez-Espigares, I., *et al.* Synthesis, molecular modelling studies and biological evaluation of new oxoeicosanoid receptor 1 agonists. *Bioorg. Med. Chem.* **26(12)**, 3580-3587 (2018).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 11/13/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM