

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



13(Z)-Eicosenoic Acid

Item No. 26169

CAS Registry No.: 17735-94-3

Synonyms: C20:1(13Z), C20:1 n-7,

cis-13-Eicosenoic Acid, Paullinic Acid

MF: $C_{20}H_{38}O_{2}$ FW: 310.5 **Purity:** ≥95% Supplied as: A neat solid Storage: -20°C Stability: ≥1 year

COOH

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

13(Z)-Eicosenoic acid is supplied as a neat solid. A stock solution may be made by dissolving the 13(Z)-eicosenoic acid in the solvent of choice. 13(Z)-Eicosenoic acid is soluble in the organic solvent ethanol, which should be purged with an inert gas, at a concentration of approximately 20 mg/ml.

13(Z)-Eicosenoic acid is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 13(Z)-eicosenoic acid should first be dissolved in ethanol and then diluted with the agueous buffer of choice. 13(Z)-Eicosenoic acid has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

13(Z)-Eicosenoic acid is an ω -7 fatty acid found in a variety of fish from the Indian, Atlantic, and Pacific oceans. It increases triglyceride accumulation in 3T3-L1 cells when used at a concentration of 50 μ M.

References

- 1. Senarath, S., Yoshinaga, K., Nagai, T., et al. Quantitative analysis of the distribution of cis-eicosenoic acid positional isomers in marine fishes from the Indian ocean. J. Oleo. Sci. 66(2), 187-197 (2017).
- 2. Senarath, S., Yoshinaga, K., Nagai, T., et al. Differential effect of cis-eicosenoic acid positional isomers on adipogenesis and lipid accumulation in 3T3-L1 cells. Eur. J. Lipid Sci. Technol. 120(6), 1700512 (2018).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 03/29/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM