

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



10(E),12(Z)-Conjugated Linoleic Acid methyl ester-d₇

Item No. 26483

Formal Name: 10E,12Z-octadecadienoic acid-16,16,17,17,18,18,18-d₇,

methyl ester

Synonyms: 10E,12Z-CLA methyl ester-d₇,

Methyl 10(E),12(Z)-Octadecadienoate methyl ester-d₇

MF: $C_{19}H_{27}D_7O_2$

301.5 FW:

Chemical Purity: ≥95% (10(E),12(Z)-Conjugated Linoleic Acid methyl ester)


Deuterium

 \geq 99% deuterated forms (d₁-d₇); \leq 1% d₀ Incorporation:

 λ_{max} : 233 nm UV/Vis.: A solution in ethanol Supplied as:

Storage: -20°C Stability: ≥1 year

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

10(E),12(Z)-Conjugated linoleic acid methyl ester-d7 is intended for use as an internal standard for the quantification of 10(E),12(Z)-conjugated linoleic acid methyl ester (Item No. 24584) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

10(E), 12(Z)-Conjugated linoleic acid methyl ester- d_7 is supplied as a solution in ethanol. To change the solvent, simply evaporate the ethanol under a gentle stream of nitrogen and immediately add the solvent of choice. Solvents such as DMSO and dimethyl formamide purged with an inert gas can be used. The solubility of 10(E),12(Z)-conjugated linoleic acid methyl ester-d₇ in these solvents is approximately 100 mg/ml.

Description

10(E),12(Z)-Conjugated linoleic acid methyl ester is a linoleic acid methyl ester containing conjugated double bonds at the 10 and 12 positions.

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 02/11/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA **PHONE:** [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM