



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



Myristoyl-L-carnitine-d₃ (chloride)

Item No. 26581

CAS Registry No.: 1334532-25-0
Formal Name: (R)-3-carboxy-N,N-dimethyl-N-(methyl-d₃)-2-(tetradecanoyloxy)propan-1-aminium, monochloride
Synonyms: L-Myristoylcarnitine-d₃, Tetradecanoyl-L-carnitine-d₃
MF: C₂₁H₃₉D₃NO₄ • Cl
FW: 411.0
Chemical Purity: ≥98% (Myristoyl-L-carnitine)
Deuterium Incorporation: ≥99% deuterated forms (d₁-d₃); ≤1% d₀
Supplied as: A solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Myristoyl-L-carnitine-d₃ (chloride) is intended for use as an internal standard for the quantification of myristoyl-L-carnitine (Item No. 26559) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

Myristoyl-L-carnitine-d₃ (chloride) is supplied as a solid. A stock solution may be made by dissolving the myristoyl-L-carnitine-d₃ (chloride) in the solvent of choice. Myristoyl-L-carnitine-d₃ (chloride) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF), which should be purged with an inert gas. The solubility of myristoyl-L-carnitine-d₃ (chloride) in ethanol and DMF is approximately 20 mg/ml and approximately 14 mg/ml in DMSO.

Description

Myristoyl-L-carnitine is a naturally occurring long-chain acylcarnitine.¹ Plasma levels of myristoyl-L-carnitine are decreased in patients with chronic fatigue syndrome and increased in patients with end-stage renal disease.^{1,2}

References

1. Reuter, S.E., Evans, A.M., Faull, R.J., *et al.* Impact of haemodialysis on individual endogenous plasma acylcarnitine concentrations in end-stage renal disease. *Ann. Clin. Biochem.* **42(Pt 5)**, 387-393 (2005).
2. Reuter, S.E. and Evans, A.M. Long-chain acylcarnitine deficiency in patients with chronic fatigue syndrome. Potential involvement of altered carnitine palmitoyltransferase-I activity. *J. Intern. Med.* **270(1)**, 76-84 (2011).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 12/11/2018

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD

ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM

WWW.CAYMANCHEM.COM