



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


PRODUCT INFORMATION



HAPC-Chol

Item No. 26584

Formal Name: (3 β)-cholest-5-en-3-ol 3-[N-[3-[(2-hydroxyethyl)amino]propyl]carbamate], hydrogen iodide
MF: C₃₃H₅₈N₂O₃ • HI
FW: 658.7
Purity: ≥95%
Supplied as: A crystalline solid
Storage: -20°C
Stability: ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

HAPC-Chol is supplied as a crystalline solid. A stock solution may be made by dissolving the HAPC-chol in the solvent of choice. HAPC-Chol is soluble in organic solvents such as ethanol and dimethyl formamide, which should be purged with an inert gas. The solubility of HAPC-chol in these solvents is approximately 10 mg/ml.

HAPC-Chol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, HAPC-chol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. HAPC-Chol has a solubility of approximately 0.14 mg/ml in a 1:6 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

HAPC-Chol is a cationic cholesterol.¹ HAPC-Chol, as part of a lipoplex with DOPE (Item No. 15091), has been used for siRNA delivery and gene silencing in MCF-7 cells in a luciferase assay without affecting cell viability. It has also been used to deliver siRNA into mice *via* intravenous injection, resulting in HAPC-chol accumulation in the lungs.

Reference

1. Hattori, Y., Nakamura, M., Takeuchi, N., *et al.* Effect of cationic lipid in cationic liposomes on siRNA delivery into the lung by intravenous injection of cationic lipoplex. *J. Drug. Target* **27(2)**, 217-227 (2019).

WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 04/05/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA

PHONE: [800] 364-9897
[734] 971-3335

FAX: [734] 971-3640

CUSTSERV@CAYMANCHEM.COM
WWW.CAYMANCHEM.COM