SZABO SCANDIC

Part of Europa Biosite

Produktinformation

- Forschungsprodukte & Biochemikalien
- Zellkultur & Verbrauchsmaterial
- Diagnostik & molekulare Diagnostik
- Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!

Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic

PRODUCT INFORMATION

OH-C-Chol

Item No. 26587

CAS Registry No.:	496801-51-5
Formal Name:	(3β)-cholest-5-en-3-ol 3-[N-[2-[(2-hydroxyethyl)amino]ethyl]carbamate]
MF:	$C_{32}H_{56}N_2O_3$
FW:	516.8
Purity:	≥95%
Supplied as:	A crystalline solid
Storage:	-20°C
Stability:	≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

OH-C-Chol is supplied as a crystalline solid. A stock solution may be made by dissolving the OH-C-chol in the solvent of choice. OH-C-Chol is soluble in organic solvents such as ethanol and dimethyl formamide, which should be purged with an inert gas. The solubility of OH-C-chol in these solvents is approximately 10 mg/ml.

OH-C-Chol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, OH-C-chol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. OH-C-Chol has a solubility of approximately 0.14 mg/ml in a 1:6 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

OH-C-Chol is a cationic cholesterol derivative.[1] OH-C-Chol, as a component of lipoplexes with DOPE (Item No. 15091), has been used for siRNA delivery and gene silencing in MCF-7 cells as well as in mice *via* intravenous injection, resulting in lipoplex accumulation in the liver.

Reference

1. Hattori, Y., Nakamura, M., Takeuchi, N., *et al.* Effect of cationic lipid in cationic liposomes on siRNA delivery into the lung by intravenous injection of cationic lipoplex. *J. Drug. Target* **27(2)**, 217-227 (2019).

CAYMAN CHEMICAL
1180 EAST ELLSWORTH RD
ANN ARBOR, MI 48108 · USA
PHONE: [800] 364-9897
[734] 971-3335
FAX: [734] 971-3640
CUSTSERV@CAYMANCHEM.COM
WWW.**CAYMANCHEM**.COM