

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



2S-Eriodictyol

Item No. 26803

CAS Registry No.: 552-58-9

Formal Name: (2S)-2-(3,4-dihydroxyphenyl)-2,3-dihydro-


5,7-dihydroxy-4H-1-benzopyran-4-one

Synonym: (+)-Eriodictyol MF: C₁₅H₁₂O₆ 288.3 FW: ≥98% **Purity:** UV/Vis.: λ_{max} : 289 nm A crystalline solid Supplied as:

Storage: -20°C Stability: ≥2 years

Plant/Unidentified Item Origin:

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

2S-Eriodictyol is supplied as a crystalline solid. A stock solution may be made by dissolving the 2S-eriodictyol in the solvent of choice, which should be purged with an inert gas. 2S-Eriodictyol is soluble in organic solvents such as DMSO and dimethyl formamide. The solubility of 2S-eriodictyol in these solvents is approximately 30 mg/ml.

2S-Eriodictyol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 2S-eriodictyol should first be dissolved in DMSO and then diluted with the aqueous buffer of choice. 2S-Eriodictyol has a solubility of approximately 0.12 mg/ml in a 1:7 solution of DMSO:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

2S-Eriodictyol a flavanone that has been found in Citrus and has antioxidant activity. It is protective against hydrogen peroxide-induced cell injury and apoptosis in EA.hy926 cells when used at a concentration of 20 μM.

Reference

1. Li, H., Li, C., Shen, T., et al. R-eriodictyol and S-eriodictyol exhibited comparable effect against H₂O₂-induced oxidative stress in EA.hy926 cells. Drug Discov. Ther. 8(5), 218-224 (2014).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

uyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 11/10/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM