



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## 1,3-Dioleoyl-2-Myristoyl Glycerol

Item No. 26872

**CAS Registry No.:** 158298-95-4  
**Formal Name:** 9Z-octadecenoic acid, 1,1'-[2-[(1-oxotetradecyl)oxy]-1,3-propanediyl] ester  
**Synonyms:** 1,3-Olein-2-Myristin, TG(18:1/14:0/18:1)  
**MF:** C<sub>53</sub>H<sub>98</sub>O<sub>6</sub>  
**FW:** 831.3  
**Purity:** ≥98%  
**Supplied as:** An oil  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

1,3-Dioleoyl-2-myristoyl glycerol is supplied as an oil. A stock solution may be made by dissolving the 1,3-dioleoyl-2-myristoyl glycerol in the solvent of choice, which should be purged with an inert gas. 1,3-dioleoyl-2-myristoyl glycerol is slightly soluble in methanol.

1,3-Dioleoyl-2-myristoyl glycerol is slightly soluble in aqueous solutions. To enhance aqueous solubility, dilute the organic solvent solution into aqueous buffers or isotonic saline. If performing biological experiments, ensure the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. We do not recommend storing the aqueous solution for more than one day.

### Description

1,3-Dioleoyl-2-myristoyl glycerol is a triacylglycerol that contains oleic acid (Item Nos. 90260 | 24659) at the *sn*-1 and *sn*-3 positions and myristic acid (Item No. 13351) at the *sn*-2 position. It has been found in palm oil.<sup>1</sup>

### Reference

1. Kallio, H., Yli-Jokipii, K., Kurvinen, J.-P., *et al.* Regioisomerism of triacylglycerols in lard, tallow, yolk, chicken skin, palm oil, palm olein, palm stearin, and a transesterified blend of palm stearin and coconut oil analyzed by tandem mass spectrometry. *J. Agric. Food Chem.* **49(7)**, 3363-3369 (2001).

**WARNING**  
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

**SAFETY DATA**  
This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the [complete](#) Safety Data Sheet, which has been sent via email to your institution.

**WARRANTY AND LIMITATION OF REMEDY**  
Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/04/2020

### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM