

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

PRODUCT INFORMATION



1-Palmitoyl-2-Lauroyl-rac-glycerol

Item No. 26875

CAS Registry No.: Formal Name:	177717-58-7 hexadecanoic acid, 3-hydroxy-2-[(1-	0
	oxododecyl)oxy]propyl ester	
Synonyms:	DG(16:0/12:0/0:0), 1-Palmitin-2-Laurin	
MF:	C ₃₁ H ₆₀ O ₅	
FW:	512.8	
Purity:	≥98%	
Supplied as:	A solid	\backslash
Storage:	-20°C	ЮН
Stability:	≥2 years	
Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis		

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

1-Palmitoyl-2-lauroyl-rac-glycerol is supplied as a solid. A stock solution may be made by dissolving the 1-palmitoyl-2-lauroyl-rac-glycerol in the solvent of choice, which should be purged with an inert gas. 1-Palmitoyl-2-lauroyl-rac-glycerol is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 1-palmitoyl-2-lauroyl-rac-glycerol in these solvents is approximately 0.25, 30, and 20 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 1-palmitoyl-2-lauroyl-rac-glycerol can be prepared by directly dissolving the solid in aqueous buffers. The solubility of 1-palmitoyl-2-lauroyl-rac-glycerol in PBS, pH 7.2, is approximately 0.7 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

1-Palmitoyl-2-lauroyl-rac-glycerol is a diacylglycerol that contains palmitic acid (Item No. 10006627) at the sn-1 position and lauric acid (Item No. 10006626) at the sn-2 position.

WARNING THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

SAFFTY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Suyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 10/03/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897 [734] 971-3335 FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.CAYMANCHEM.COM