

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# PRODUCT INFORMATION



## 1,3-Dipalmitoyl Glycerol

Item No. 26974

CAS Registry No.: 502-52-3

Formal Name: hexadecanoic acid, 1,1'-(2-hydroxy-

1,3-propanediyl) ester

Synonyms: DG(16:0/0:0/16:0), 1,3-Dipalmitin

MF:  $C_{35}H_{68}O_{5}$ FW: 568.9 **Purity:** ≥98% Supplied as: A solid Storage: -20°C Stability: ≥2 years

ОН

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### **Laboratory Procedures**

1,3-Dipalmitoyl glycerol is supplied as a solid. A stock solution may be made by dissolving the 1,3-dipalmitoyl glycerol in the solvent of choice, which should be purged with an inert gas. 1,3-Dipalmitoyl glycerol is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide. The solubility of 1,3-dipalmitoyl glycerol in these solvents is approximately 0.25, 30, and 20 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of 1,3-dipalmitoyl glycerol can be prepared by directly dissolving the solid in aqueous buffers. The solubility of 1,3-dipalmitoyl glycerol in PBS, pH 7.2, is approximately 0.7 mg/ml. We do not recommend storing the aqueous solution for more than one day.

#### Description

1,3-Dipalmitoyl glycerol is a diacylglycerol that contains palmitic acid (Item No. 10006627) at the sn-1 and sn-3 positions. It activates protein kinase C  $\alpha$  (PKC $\alpha$ ) in vitro (K<sub>a</sub> = 3.8  $\mu$ M). 1,3-Dipalmitoyl glycerol (0.25-2 µM) inhibits apoptosis and production of reactive oxygen species (ROS) and pro-inflammatory mediators induced by oxygen-glucose deprivation and reperfusion (OGD/R) in SH-SY5Y neuroblastoma cells.2

#### References

- 1. Sánchez-Piñera, P., Micol, V., Corbalán-García, S., et al. A comparative study of the activation of protein kinase C α by different diacylglycerol isomers. Biochem. J. 337(3), 387-395 (1999).
- 2. Cheng, M.-C. and Pan, T.-M. Glyceryl 1,3-dipalmitate produced from Lactobacillus paracasei subspecies. paracasei NTU 101 inhibits oxygen-glucose deprivation and reperfusion-induced oxidative stress via upregulation of peroxisome proliferator-activated receptor γ in neuronal SH-SY5Y cells. J. Agric. Food Chem. 65(36), 7926-7933 (2017).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

## WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 06/09/2020

### **CAYMAN CHEMICAL**

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM