

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



C29

Item No. 27029

CAS Registry No.: 363600-92-4

Formal Name: 3-[[(2-hydroxy-3-methoxyphenyl)

methylenelaminol-2-methyl-


benzoic acid

Synonym: TLR2-IN-C29 MF: C₁₆H₁₅NO₄ FW: 285.3 **Purity:**

UV/Vis.: λ_{max} : 226, 277, 320 nm Supplied as: A crystalline solid

-20°C Storage: Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

C29 is supplied as a crystalline solid. A stock solution may be made by dissolving the C29 in the solvent of choice. C29 is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of C29 in these solvents is approximately 0.5, 11, and 10 mg/ml, respectively.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of C29 can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of C29 in PBS, pH 7.2, is approximately 0.3 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

C29 is an inhibitor of toll-like receptor 2 (TLR2) signaling.¹ It inhibits synthetic and bacterial TLR2 agonist-induced TLR2/1 and TLR2/6 signaling in HEK-293T cells expressing human TLR2 (HEK-TLR2) and THP-1 cells when used at a concentration of 50 μM. C29 also inhibits TLR2/1 signaling in primary murine macrophages without inducing cytotoxicity. It decreases the expression of IL-8 induced by the synthetic bacterial lipopeptide TLR2 agonist Pam₃CSK₄ (Item No. 24126) in HEK-TLR2 cells. C29 (150 μM) inhibits the Pam₂CSK₄-induced interaction between TLR2 and the adapter protein MyD88, blocks MAPK activation, and decreases NF-κB activation and IκBα degradation in THP-1 cells.

Reference

1. Mistry, P., Laird, M.H., Schwarz, R.S., et al. Inhibition of TLR2 signaling by small molecule inhibitors targeting a pocket within the TLR2 TIR domain. Proc. Natl. Acad. Sci. U.S.A. 112(17), 5455-5460 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the material can be found on our website.

Copyright Cayman Chemical Company, 06/05/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM