

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



β-Muricholic Acid-d₄

Item No. 27035

 (5β) - 3α , 6β , 7β -trihydroxy-cholan-2,2,4,4-d₄-Formal Name:

24-oic acid

5β-Cholanic Acid-3α,6β,7β-triol-d₄, Synonyms:

β-MCA-d₄

MF: $C_{24}H_{36}D_4O_5$ FW: 412.6

Chemical Purity: ≥98% (β-Muricholic Acid)


Deuterium

Incorporation: \geq 99% deuterated forms (d₁-d₄); \leq 1% d₀

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.



Laboratory Procedures

 β -Muricholic acid-d₄ (β -MCA-d₄) is intended for use as an internal standard for the quantification of β-muricholic acid (Item No. 20287) by GC- or LC-MS. The accuracy of the sample weight in this vial is between 5% over and 2% under the amount shown on the vial. If better precision is required, the deuterated standard should be quantitated against a more precisely weighed unlabeled standard by constructing a standard curve of peak intensity ratios (deuterated versus unlabeled).

 β -MCA- d_4 is supplied as a crystalline solid. A stock solution may be made by dissolving the β -MCA- d_4 in the solvent of choice, which should be purged with an inert gas. β -MCA-d₄ is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of β-MCA-d₄ in ethanol and DMSO is approximately 20 mg/ml and approximately 30 mg/ml in DMF.

Description

β-MCA is a murine-specific primary bile acid.^{1,2} Dietary administration of β-MCA reduces HMG-CoA reductase activity in liver microsomes from mice fed a high cholesterol and cholic acid diet.³ Dietary administration of β-MCA also dissolves 100% of gallstones in a gallstone-susceptible mouse model of diet-induced cholesterol gallstones.4

References

- 1. Eyssen, H.J., Parmentier, G.G., and Mertens, J.A. Sulfate bile acids in germ-free and conventional mice. Eur. J. Biochem. 66(3), 507-514 (1976).
- 2. Wahlström, A., Sayin, S.I., Marschall, H.-I., et al. Intestinal crosstalk between bile acids and microbiota and its impact on host metabolism. Cell Metab. 24(1), 41-50 (2016).
- 3. Fujino, Y., Nakayama, K., Yoshimura, K., et al. Suppression of hepatic HMG-CoA reductase activity by β-muricholic acid in mice fed a diet containing cholesterol and cholic acid. Jpn. J. Pharmacol. 46(4), 421-423 (1988).
- 4. Wang, D.Q.-H. and Tazuma, S. Effect of β-muricholic acid on the prevention and dissolution of cholesterol gallstones in C57L/J mice. J. Lipid. Res. 43(11), 1960-1968 (2002).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information Buyer agrees to purchase the mater can be found on our website.

Copyright Cayman Chemical Company, 05/27/2021

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM