



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 


# PRODUCT INFORMATION



## 1,2-Dilinoleoyl-3-Stearoyl-*rac*-glycerol

Item No. 27063

**CAS Registry No.:** 2190-17-2  
**Formal Name:** 9Z,12Z-octadecadienoic acid, 1,1'-[1-[[[(1-oxooctadecyl)oxy] methyl]-1,2-ethanediyl] ester  
**Synonyms:** 1,2-Linolein-3-Stearin, TG(18:2/18:2/18:0)  
**MF:** C<sub>57</sub>H<sub>102</sub>O<sub>6</sub>  
**FW:** 883.4  
**Purity:** ≥98%  
**Supplied as:** A liquid  
**Storage:** -20°C  
**Stability:** ≥2 years



Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

### Laboratory Procedures

1,2-Dilinoleoyl-3-stearoyl-*rac*-glycerol is supplied as a liquid. A stock solution may be made by dissolving the 1,2-dilinoleoyl-3-stearoyl-*rac*-glycerol in the solvent of choice. 1,2-Dilinoleoyl-3-stearoyl-*rac*-glycerol is soluble in organic solvents such as ethanol and dimethyl formamide, which should be purged with an inert gas. The solubility of 1,2-dilinoleoyl-3-stearoyl-*rac*-glycerol in these solvents is approximately 10 mg/ml. 1,2-Dilinoleoyl-3-stearoyl-*rac*-glycerol is also slightly soluble in chloroform.

1,2-Dilinoleoyl-3-stearoyl-*rac*-glycerol is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, 1,2-dilinoleoyl-3-stearoyl-*rac*-glycerol should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. 1,2-Dilinoleoyl-3-stearoyl-*rac*-glycerol has a solubility of approximately 0.5 mg/ml in a 1:1 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

### Description

1,2-Dilinoleoyl-3-stearoyl-*rac*-glycerol is a triacylglycerol that contains linoleic acid (Item Nos. 90150 | 90150.1 | 21909) at the *sn*-1 and *sn*-2 positions and stearic acid (Item No. 10011298) at the *sn*-3 position. It has been found in various plant oils, including soybean, poppy seed, maize, evening primrose, and blackcurrant oils.<sup>1</sup>

### Reference

1. Mottram, H.R., Woodbury, S.E., and Evershed, R.P. Identification of triacylglycerol positional isomers present in vegetable oils by high performance liquid chromatography/atmospheric pressure chemical ionization mass spectrometry. *Rapid Commun. Mass Spectrom.* **11(12)**, 1240-1252 (1997).

#### WARNING

THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

#### SAFETY DATA

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

#### WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 05/29/2019

#### CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD  
ANN ARBOR, MI 48108 · USA

**PHONE:** [800] 364-9897  
[734] 971-3335

**FAX:** [734] 971-3640

CUSTSERV@CAYMANCHEM.COM  
WWW.CAYMANCHEM.COM