

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



PAR4 (1-6) (human)

Item No. 27126

CAS Registry No.: 225779-44-2

Formal Name: glycyl-L-tyrosyl-L-prolylglycyl-L-

glutaminyl-L-valine

Synonym: **GYPGQV**

MF: $C_{28}H_{41}N_7O_9$

619.7 FW: ≥98% **Purity:**

UV/Vis.: λ_{max} : 225, 279 nm Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

H-Gly-Tyr-Pro-Gly-Gln-Val-OH

Laboratory Procedures

PAR4 (1-6) (human) is supplied as a crystalline solid. A stock solution may be made by dissolving the PAR4 (1-6) (human) in the solvent of choice. PAR4 (1-6) (human) is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide, which should be purged with an inert gas. The solubility of PAR4 (1-6) (human) in these solvents is approximately 50 mg/ml.

Further dilutions of the stock solution into aqueous buffers or isotonic saline should be made prior to performing biological experiments. Ensure that the residual amount of organic solvent is insignificant, since organic solvents may have physiological effects at low concentrations. Organic solvent-free aqueous solutions of PAR4 (1-6) (human) can be prepared by directly dissolving the crystalline solid in aqueous buffers. The solubility of PAR4 (1-6) (human) in PBS, pH 7.2, is approximately 10 mg/ml. We do not recommend storing the aqueous solution for more than one day.

Description

PAR4 (1-6) is a peptide agonist of proteinase-activated receptor 4 (PAR4) that corresponds to residues 1-6 of the amino terminal tethered ligand sequence of human PAR4 and residues 48-53 of the full-length sequence.¹ It activates PAR4 and the cleavage site mutant PAR4^{R47A} when used at a concentration of 500 μM.² PAR4 (1-6) induces platelet aggregation of isolated washed human platelets when used at a concentration of 1 mM but does not affect clotting time induced by factor VIIa, soluble tissue factor, and collagen in an ex vivo coagulation assay.3

References

- 1. Vergnolle, N., Wallace, J.L., Bunnett, N.W., et al. Protease-activated receptors in inflammation, neuronal signaling and pain. Trends Pharmacol. Sci. 22(3), 146-152 (2001).
- 2. Xu, W.F., Andersen, H., Whitmore, T.E., et al. Cloning and characterization of human protease-activated receptor 4. Proc. Natl. Acad. Sci. U.S.A. 95(12), 6642-6646 (1998).
- 3. Andersen, H., Greenberg, D.L., Fujikawa, K., et al. Protease-activated receptor 1 is the primary mediator of thrombin-stimulated platelet procoagulant activity. Proc. Natl. Acad. Sci. U.S.A. 96(20), 11189-11193 (1999).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

Buyer agrees to purchase the material subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website.

Copyright Cayman Chemical Company, 06/18/2019

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM