

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



PRODUCT INFORMATION



Oleanolic Acid Acrylate

Item No. 27206

CAS Registry No.: 1975976-24-9

Formal Name: 3β -[(1-oxo-2-propen-1-yl)oxy]-

olean-12-en-28-oic acid

Synonym: MF: $C_{33}H_{50}O_4$ FW: 510.7 **Purity:** ≥95%

Supplied as: A crystalline solid

Storage: -20°C Stability: ≥2 years

Information represents the product specifications. Batch specific analytical results are provided on each certificate of analysis.

Laboratory Procedures

Oleanolic acid acrylate (OAA) is supplied as a crystalline solid. A stock solution may be made by dissolving the OAA in the solvent of choice, which should be purged with an inert gas. OAA is soluble in organic solvents such as ethanol, DMSO, and dimethyl formamide (DMF). The solubility of OAA in ethanol and DMF is approximately 30 mg/ml and approximately 20 mg/ml in DMSO.

OAA is sparingly soluble in aqueous buffers. For maximum solubility in aqueous buffers, OAA should first be dissolved in ethanol and then diluted with the aqueous buffer of choice. OAA has a solubility of approximately 0.02 mg/ml in a 1:40 solution of ethanol:PBS (pH 7.2) using this method. We do not recommend storing the aqueous solution for more than one day.

Description

OAA is an agonist of the serotonin (5-HT) receptor subtype $5\text{-HT}_{1\text{A}}$ and a derivative of oleanolic acid (Item No. 11726). It acts as an agonist at the 5-HT $_{1A}$ receptor in mouse hippocampal homogenates in a [35 S]GTP γ S binding assay (EC $_{50}$ = 0.79 μ M). It is also a selective inhibitor of MAO-A over MAO-B (IC₅₀s = 48.8μ M and >100 μ M, respectively). In mice, OAA (10 mg/kg) reduces immobility in the forced swim test and increases the number of crossings and time spent at the center in the open field test, indicating anti-depressant-like and anxiolytic activities.

Reference

1. Fajemiroye, J.O., Polepally, P.R., Chaurasiya, N.D., et al. Oleanolic acid acrylate elicits antidepressant-like effect mediated by 5-HT_{1A} receptor. Sci. Rep. 5, 11582 (2015).

WARNING
THIS PRODUCT IS FOR RESEARCH ONLY - NOT FOR HUMAN OR VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

This material should be considered hazardous until further information becomes available. Do not ingest, inhale, get in eyes, on skin, or on clothing. Wash thoroughly after handling. Before use, the user must review the complete Safety Data Sheet, which has been sent via email to your institution.

WARRANTY AND LIMITATION OF REMEDY

subject to Cayman's Terms and Conditions. Complete Terms and Conditions including Warranty and Limitation of Liability information can be found on our website

Copyright Cayman Chemical Company, 01/28/2020

CAYMAN CHEMICAL

1180 EAST ELLSWORTH RD ANN ARBOR, MI 48108 · USA PHONE: [800] 364-9897

[734] 971-3335

FAX: [734] 971-3640 CUSTSERV@CAYMANCHEM.COM WWW.**CAYMANCHEM**.COM